

# Irrationality Criteria for Series by Erdős and Straus

Angeliki Koutsoukou-Argyraki and Wenda Li

February 6, 2026

## Abstract

We formalise certain irrationality criteria for infinite series of the form:

$$\sum_n \frac{b_n}{\prod_{i \leq n} a_i}$$

where  $b_n, a_i$  are integers. The result is due to P. Erdős and E.G. Straus [1], and in particular we formalise Theorem 2.1, Corollary 2.10 and Theorem 3.1. The latter is an application of Theorem 2.1 involving the prime numbers.

## Contents

<b>1</b>	<b>Miscellaneous</b>	<b>1</b>
<b>2</b>	<b>Theorem 2.1 and Corollary 2.10</b>	<b>2</b>
<b>3</b>	<b>Some auxiliary results on the prime numbers.</b>	<b>3</b>
<b>4</b>	<b>Theorem 3.1</b>	<b>4</b>
<b>5</b>	<b>Acknowledgements</b>	<b>4</b>

**theory** *Irrational-Series-Erdos-Straus* **imports**  
*Prime-Number-Theorem.Prime-Number-Theorem*  
*Prime-Distribution-Elementary.PNT-Consequences*  
**begin**

## 1 Miscellaneous

**lemma** *suminf-comparison*:  
**assumes** *summable f* **and** *gf:  $\bigwedge n. \text{norm } (g\ n) \leq f\ n$*   
**shows** *suminf g  $\leq$  suminf f*  
{*proof*}

**lemma** *tendsto-of-int-diff-0*:  
**assumes**  $(\lambda n. f\ n - \text{of-int}(g\ n)) \longrightarrow (0::\text{real}) \forall_F n$  in sequentially.  $f\ n > 0$   
**shows**  $\forall_F n$  in sequentially.  $0 \leq g\ n$   
 $\langle \text{proof} \rangle$

**lemma** *eventually-mono-sequentially*:  
**assumes** eventually  $P$  sequentially  
**assumes**  $\bigwedge x. P\ (x+k) \implies Q\ (x+k)$   
**shows** eventually  $Q$  sequentially  
 $\langle \text{proof} \rangle$

**lemma** *frequently-eventually-at-top*:  
**fixes**  $P\ Q::'a::\text{linorder} \Rightarrow \text{bool}$   
**assumes** frequently  $P$  at-top eventually  $Q$  at-top  
**shows** frequently  $(\lambda x. P\ x \wedge (\forall y \geq x. Q\ y))$  at-top  
 $\langle \text{proof} \rangle$

**lemma** *eventually-at-top-mono*:  
**fixes**  $P\ Q::'a::\text{linorder} \Rightarrow \text{bool}$   
**assumes** event- $P$ :eventually  $P$  at-top  
**assumes**  $PQ\text{-imp}:\bigwedge x. x \geq z \implies \forall y \geq x. P\ y \implies Q\ x$   
**shows** eventually  $Q$  at-top  
 $\langle \text{proof} \rangle$

**lemma** *frequently-at-top-elim*:  
**fixes**  $P\ Q::'a::\text{linorder} \Rightarrow \text{bool}$   
**assumes**  $\exists_F x$  in at-top.  $P\ x$   
**assumes**  $\bigwedge i. P\ i \implies \exists j > i. Q\ j$   
**shows**  $\exists_F x$  in at-top.  $Q\ x$   
 $\langle \text{proof} \rangle$

**lemma** *less-Liminf-iff*:  
**fixes**  $X :: - \Rightarrow - :: \text{complete-linorder}$   
**shows**  $\text{Liminf } F\ X < C \iff (\exists y < C. \text{frequently } (\lambda x. y \geq X\ x)\ F)$   
 $\langle \text{proof} \rangle$

**lemma** *sequentially-even-odd-imp*:  
**assumes**  $\forall_F N$  in sequentially.  $P\ (2*N) \forall_F N$  in sequentially.  $P\ (2*N+1)$   
**shows**  $\forall_F n$  in sequentially.  $P\ n$   
 $\langle \text{proof} \rangle$

## 2 Theorem 2.1 and Corollary 2.10

**context**  
**fixes**  $a\ b :: \text{nat} \Rightarrow \text{int}$   
**assumes**  $a\text{-pos}$ :  $\forall n. a\ n > 0$  **and**  $a\text{-large}$ :  $\forall_F n$  in sequentially.  $a\ n > 1$   
**and**  $ab\text{-tendsto}$ :  $(\lambda n. |b\ n| / (a\ (n-1) * a\ n)) \longrightarrow 0$   
**begin**

**private lemma** *aux-series-summable*: *summable*  $(\lambda n. b\ n / (\prod_{k \leq n} a\ k))$   
 $\langle \text{proof} \rangle$  **fun** *get-c*:: $(\text{nat} \Rightarrow \text{int}) \Rightarrow (\text{nat} \Rightarrow \text{int}) \Rightarrow \text{int} \Rightarrow \text{nat} \Rightarrow (\text{nat} \Rightarrow \text{int})$  **where**  
*get-c*  $a' b' B\ N\ 0 = \text{round } (B * b' N / a' N)$   
*get-c*  $a' b' B\ N\ (\text{Suc } n) = \text{get-c } a' b' B\ N\ n * a' (n+N) - B * b' (n+N)$

**lemma** *ab-rationality-imp*:

**assumes** *ab-rational*: $(\sum n. (b\ n / (\prod_{i \leq n} a\ i))) \in \mathbb{Q}$

**shows**  $\exists (B::\text{int}) > 0. \exists c::\text{nat} \Rightarrow \text{int}.$

$(\forall_F n \text{ in sequentially. } B * b\ n = c\ n * a\ n - c(n+1) \wedge |c(n+1)| < a\ n/2)$

$\wedge (\lambda n. c\ (\text{Suc } n) / a\ n) \longrightarrow 0$

$\langle \text{proof} \rangle$  **lemma** *imp-ab-rational*:

**assumes**  $\exists (B::\text{int}) > 0. \exists c::\text{nat} \Rightarrow \text{int}.$

$(\forall_F n \text{ in sequentially. } B * b\ n = c\ n * a\ n - c(n+1) \wedge |c(n+1)| < a$

$n/2)$

**shows**  $(\sum n. (b\ n / (\prod_{i \leq n} a\ i))) \in \mathbb{Q}$

$\langle \text{proof} \rangle$

**theorem** *theorem-2-1-Erdos-Straus* :

$(\sum n. (b\ n / (\prod_{i \leq n} a\ i))) \in \mathbb{Q} \longleftrightarrow (\exists (B::\text{int}) > 0. \exists c::\text{nat} \Rightarrow \text{int}.$

$(\forall_F n \text{ in sequentially. } B * b\ n = c\ n * a\ n - c(n+1) \wedge |c(n+1)| < a\ n/2))$

$\langle \text{proof} \rangle$

The following is a Corollary to Theorem 2.1.

**corollary** *corollary-2-10-Erdos-Straus*:

**assumes** *ab-event*: $\forall_F n \text{ in sequentially. } b\ n > 0 \wedge a\ (n+1) \geq a\ n$

**and** *ba-lim-leq*: $\lim (\lambda n. (b(n+1) - b\ n) / a\ n) \leq 0$

**and** *ba-lim-exist*:*convergent*  $(\lambda n. (b(n+1) - b\ n) / a\ n)$

**and** *liminf*  $(\lambda n. a\ n / b\ n) = 0$

**shows**  $(\sum n. (b\ n / (\prod_{i \leq n} a\ i))) \notin \mathbb{Q}$

$\langle \text{proof} \rangle$

**end**

### 3 Some auxiliary results on the prime numbers.

**lemma** *nth-prime-nonzero[simp]*:*nth-prime*  $n \neq 0$

$\langle \text{proof} \rangle$

**lemma** *nth-prime-gt-zero[simp]*:*nth-prime*  $n > 0$

$\langle \text{proof} \rangle$

**lemma** *ratio-of-consecutive-primes*:

$(\lambda n. \text{nth-prime } (n+1) / \text{nth-prime } n) \longrightarrow 1$

$\langle \text{proof} \rangle$

**lemma** *nth-prime-double-sqrt-less*:

**assumes**  $\varepsilon > 0$

**shows**  $\forall_F n \text{ in sequentially. } (\text{nth-prime } (2*n) - \text{nth-prime } n)$

$/ \text{sqrt } (\text{nth-prime } n) < n \text{ powr } (1/2 + \varepsilon)$

*<proof>*

## 4 Theorem 3.1

Theorem 3.1 is an application of Theorem 2.1 with the sequences considered involving the prime numbers.

**theorem** *theorem-3-10-Erdos-Straus:*

**fixes**  $a::nat \Rightarrow int$

**assumes**  $a\text{-pos}:\forall n. a\ n > 0$  **and** *mono*  $a$

**and**  $nth\text{-}1:(\lambda n. nth\text{-}prime\ n / (a\ n)^2) \longrightarrow 0$

**and**  $nth\text{-}2:\liminf (\lambda n. a\ n / nth\text{-}prime\ n) = 0$

**shows**  $(\sum n. (nth\text{-}prime\ n / (\prod_{i \leq n} a\ i))) \notin \mathbb{Q}$

*<proof>*

## 5 Acknowledgements

A.K.-A. and W.L. were supported by the ERC Advanced Grant ALEXANDRIA (Project 742178) funded by the European Research Council and led by Professor Lawrence Paulson at the University of Cambridge, UK.

**end**

## References

- [1] P. Erdős and E. Straus. On the irrationality of certain series. *Pacific journal of mathematics*, 55(1):85–92, 1974.