

Real-Valued Special Functions: Upper and Lower Bounds

Lawrence C. Paulson

May 26, 2024

Abstract

This development proves upper and lower bounds for several familiar real-valued functions. For \sin , \cos , \exp and the square root function, it defines and verifies infinite families of upper and lower bounds, mostly based on Taylor series expansions. For \tan^{-1} , \ln and \exp , it verifies a finite collection of upper and lower bounds, originally obtained from the functions' continued fraction expansions using the computer algebra system Maple. A common theme in these proofs is to take the difference between a function and its approximation, which should be zero at one point, and then consider the sign of the derivative.

The immediate purpose of this development is to verify axioms used by MetiTarski [1], an automatic theorem prover for real-valued special functions. Crucial to MetiTarski's operation is the provision of upper and lower bounds for each function of interest.

Contents

1	General Lemmas for Proving Function Inequalities	3
2	Arctan Upper and Lower Bounds	5
2.1	Upper Bound 1	5
2.2	Lower Bound 1	8
2.3	Upper Bound 3	9
2.4	Lower Bound 3	12
2.5	Upper Bound 4	12
2.6	Lower Bound 4	15
3	Exp Upper and Lower Bounds	17
3.1	Taylor Series Bounds	17
3.2	Continued Fraction Bound 2	20
3.3	Continued Fraction Bound 3	21
3.4	Continued Fraction Bound 4	23
3.5	Continued Fraction Bound 5	24
3.5.1	Proving monotonicity via a non-negative derivative . .	25
3.5.2	Results	25
3.6	Continued Fraction Bound 6	26
3.7	Continued Fraction Bound 7	28
3.7.1	Proving monotonicity via a non-negative derivative . .	28
3.7.2	Results	29
4	Log Upper and Lower Bounds	31
4.1	Upper Bound 3	31
4.2	Upper Bound 5	32
4.3	Upper Bound 7	33
4.4	Upper Bound 9	34
4.5	Upper Bound 11	35
4.6	Upper Bound 13	36
4.7	Upper Bound 15	37

5	Sine and Cosine Upper and Lower Bounds	40
5.1	Simple base cases	40
5.2	Taylor series approximants	41
5.3	Inductive proof of sine inequalities	42
5.4	Collecting the results	43
6	Square Root Upper and Lower Bounds	45
6.1	Upper bounds	45
6.2	Lower bounds	47

Chapter 1

General Lemmas for Proving Function Inequalities

```
theory Bounds-Lemmas
imports Complex-Main
```

```
begin
```

These are for functions that are differentiable over a closed interval.

```
lemma gen-lower-bound-increasing:
```

```
fixes a :: real
```

```
assumes a ≤ x
```

```
and ∀y. a ≤ y ⇒ y ≤ x ⇒ ((λx. fl x - f x) has-real-derivative g y) (at y)
```

```
and ∀y. a ≤ y ⇒ y ≤ x ⇒ g y ≤ 0
```

```
and fl a = f a
```

```
shows fl x ≤ f x
```

```
proof -
```

```
have fl x - f x ≤ fl a - f a
```

```
apply (rule DERIV-nonpos-imp-nonincreasing [where f = λx. fl x - f x])
```

```
apply (rule assms)
```

```
apply (intro allI impI exI conjI)
```

```
apply (rule assms | simp)+
```

```
done
```

```
also have ... = 0
```

```
by (simp add: assms)
```

```
finally show ?thesis
```

```
by simp
```

```
qed
```

```
lemma gen-lower-bound-decreasing:
```

```
fixes a :: real
```

```
assumes x ≤ a
```

```
and ∀y. x ≤ y ⇒ y ≤ a ⇒ ((λx. fl x - f x) has-real-derivative g y) (at y)
```

```
and ∀y. x ≤ y ⇒ y ≤ a ⇒ g y ≥ 0
```

```
and fl a = f a
```

```

shows  $f l x \leq f x$ 
proof -
have  $f l (-(-x)) \leq f (-(-x))$ 
apply (rule gen-lower-bound-increasing [of  $-a - x - \lambda u. - g (-u)$ ])
apply (auto simp: assms)
apply (subst DERIV-mirror [symmetric])
apply (simp add: assms)
done
then show ?thesis
by simp
qed

lemma gen-upper-bound-increasing:
fixes  $a :: real$ 
assumes  $a \leq x$ 
and  $\bigwedge y. a \leq y \implies y \leq x \implies ((\lambda x. fu x - f x) \text{ has-real-derivative } g y) \text{ (at } y\text{)}$ 
and  $\bigwedge y. a \leq y \implies y \leq x \implies g y \geq 0$ 
and  $fu a = f a$ 
shows  $f x \leq fu x$ 
apply (rule gen-lower-bound-increasing [of  $a x f fu \lambda u. - g u$ ])
using assms DERIV-minus [where  $f = \lambda x. fu x - f x$ ]
apply auto
done

lemma gen-upper-bound-decreasing:
fixes  $a :: real$ 
assumes  $x \leq a$ 
and  $\bigwedge y. x \leq y \implies y \leq a \implies ((\lambda x. fu x - f x) \text{ has-real-derivative } g y) \text{ (at } y\text{)}$ 
and  $\bigwedge y. x \leq y \implies y \leq a \implies g y \leq 0$ 
and  $fu a = f a$ 
shows  $f x \leq fu x$ 
apply (rule gen-lower-bound-decreasing [of  $x a - \lambda u. - g u$ ])
using assms DERIV-minus [where  $f = \lambda x. fu x - f x$ ]
apply auto
done

end

```

Chapter 2

Arctan Upper and Lower Bounds

```
theory Atan-CF-Bounds
imports Bounds-Lemmas
HOL-Library.Sum-of-Squares
```

```
begin
```

Covers all bounds used in arctan-upper.ax, arctan-lower.ax and arctan-extended.ax, excepting only arctan-extended2.ax, which is used in two atan-error-analysis problems.

2.1 Upper Bound 1

```
definition arctan-upper-11 :: real ⇒ real
  where arctan-upper-11 ≡ λx. -(pi/2) - 1/x
```

```
definition diff-delta-arctan-upper-11 :: real ⇒ real
  where diff-delta-arctan-upper-11 ≡ λx. 1 / (x^2 * (1 + x^2))
```

```
lemma d-delta-arctan-upper-11: x ≠ 0 ==>
  ((λx. arctan-upper-11 x - arctan x) has-field-derivative diff-delta-arctan-upper-11
  x) (at x)
unfolding arctan-upper-11-def diff-delta-arctan-upper-11-def
apply (intro derivative-eq-intros | simp)+
apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
done
```

```
lemma d-delta-arctan-upper-11-pos: x ≠ 0 ==> diff-delta-arctan-upper-11 x > 0
unfolding diff-delta-arctan-upper-11-def
by (simp add: divide-simps zero-less-mult-iff add-pos-pos)
```

Different proof needed here: they coincide not at zero, but at (-) infinity!

```
lemma arctan-upper-11:
```

```

assumes  $x < 0$ 
shows  $\arctan(x) < \arctan\text{-upper-11 } x$ 
proof -
have  $((\lambda x. \arctan\text{-upper-11 } x - \arctan x) \longrightarrow -(\pi / 2) - 0 - (-(\pi / 2)))$ 
at-bot
  unfolding  $\arctan\text{-upper-11}\text{-def}$ 
  apply (intro tendsto-intros tendsto-arctan-at-bot, auto simp: ext [OF divide-inverse])
  apply (metis tendsto-inverse-0 at-bot-le-at-infinity tendsto-mono)
  done
then have *:  $((\lambda x. \arctan\text{-upper-11 } x - \arctan x) \longrightarrow 0)$  at-bot
  by simp
have  $0 < \arctan\text{-upper-11 } x - \arctan x$ 
  apply (rule DERIV-pos-imp-increasing-at-bot [OF - *])
  apply (metis assms d-delta-arctan-upper-11 d-delta-arctan-upper-11-pos not-le)
  done
then show ?thesis
  by auto
qed

```

```

definition  $\arctan\text{-upper-12} :: \text{real} \Rightarrow \text{real}$ 
where  $\arctan\text{-upper-12} \equiv \lambda x. 3*x / (x^2 + 3)$ 

```

```

definition  $\text{diff-delta-arctan}\text{-upper-12} :: \text{real} \Rightarrow \text{real}$ 
where  $\text{diff-delta-arctan}\text{-upper-12} \equiv \lambda x. -4*x^4 / ((x^2+3)^2 * (1+x^2))$ 

```

```

lemma  $d\text{-delta-arctan}\text{-upper-12}:$ 
 $((\lambda x. \arctan\text{-upper-12 } x - \arctan x) \text{ has-field-derivative } \text{diff-delta-arctan}\text{-upper-12}$ 
 $x) \text{ (at } x)$ 
  unfolding  $\arctan\text{-upper-12}\text{-def diff-delta-arctan}\text{-upper-12}\text{-def}$ 
  apply (intro derivative-eq-intros, simp-all)
  apply (auto simp: divide-simps add-nonneg-eq-0-iff, algebra)
  done

```

Strict inequalities also possible

```

lemma  $\arctan\text{-upper-12}:$ 
assumes  $x \leq 0$  shows  $\arctan(x) \leq \arctan\text{-upper-12 } x$ 
apply (rule gen-upper-bound-decreasing [OF assms d-delta-arctan-upper-12])
apply (auto simp: diff-delta-arctan-upper-12-def arctan-upper-12-def)
done

```

```

definition  $\arctan\text{-upper-13} :: \text{real} \Rightarrow \text{real}$ 
where  $\arctan\text{-upper-13} \equiv \lambda x. x$ 

```

```

definition  $\text{diff-delta-arctan}\text{-upper-13} :: \text{real} \Rightarrow \text{real}$ 
where  $\text{diff-delta-arctan}\text{-upper-13} \equiv \lambda x. x^2 / (1 + x^2)$ 

```

```

lemma  $d\text{-delta-arctan}\text{-upper-13}:$ 
 $((\lambda x. \arctan\text{-upper-13 } x - \arctan x) \text{ has-field-derivative } \text{diff-delta-arctan}\text{-upper-13}$ 
 $x) \text{ (at } x)$ 

```

```

unfolding arctan-upper-13-def diff-delta-arctan-upper-13-def
apply (intro derivative-eq-intros, simp-all)
apply (simp add: divide-simps add-nonneg-eq-0-iff)
done

lemma arctan-upper-13:
  assumes  $x \geq 0$  shows  $\arctan(x) \leq \text{arctan-upper-13 } x$ 
  apply (rule gen-upper-bound-increasing [OF assms d-delta-arctan-upper-13])
  apply (auto simp: diff-delta-arctan-upper-13-def arctan-upper-13-def)
  done

definition arctan-upper-14 :: real  $\Rightarrow$  real
  where arctan-upper-14  $\equiv \lambda x. \pi/2 - 3*x / (1 + 3*x^2)$ 

definition diff-delta-arctan-upper-14 :: real  $\Rightarrow$  real
  where diff-delta-arctan-upper-14  $\equiv \lambda x. -4 / ((1 + 3*x^2)^2 * (1+x^2))$ 

lemma d-delta-arctan-upper-14:
   $((\lambda x. \arctan\text{-upper-14 } x - \arctan x) \text{ has-field-derivative } \text{diff-delta-arctan-upper-14}$ 
 $x)$  (at  $x$ )
unfolding arctan-upper-14-def diff-delta-arctan-upper-14-def
apply (intro derivative-eq-intros | simp add: add-nonneg-eq-0-iff)+
apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
done

lemma d-delta-arctan-upper-14-neg: diff-delta-arctan-upper-14  $x < 0$ 
unfolding diff-delta-arctan-upper-14-def
apply (auto simp: divide-simps add-nonneg-eq-0-iff zero-less-mult-iff)
using power2-less-0 [of  $x$ ]
apply arith
done

lemma lim14:  $((\lambda x::\text{real}. 3 * x / (1 + 3 * x^2)) \xrightarrow{} 0)$  at-infinity!
apply (rule tendsto-0-le [where  $f = \text{inverse}$  and  $K=1$ ])
apply (metis tendsto-inverse-0)
apply (simp add: eventually-at-infinity)
apply (rule-tac  $x=1$  in exI)
apply (simp add: power-eq-if abs-if divide-simps add-sign-intros)
done

```

Different proof needed here: they coincide not at zero, but at (+) infinity!

```

lemma arctan-upper-14:
  assumes  $x > 0$ 
  shows  $\arctan(x) < \text{arctan-upper-14 } x$ 
proof -
  have  $((\lambda x. \arctan\text{-upper-14 } x - \arctan x) \xrightarrow{} \pi/2 - 0 - \pi/2)$  at-top
  unfolding arctan-upper-14-def
  apply (intro tendsto-intros tendsto-arctan-at-top)
  apply (auto simp: tendsto-mono [OF at-top-le-at-infinity lim14])

```

```

done
then have *:  $((\lambda x. \text{arctan-upper-14 } x - \text{arctan } x) \longrightarrow 0)$  at-top
  by simp
have  $0 < \text{arctan-upper-14 } x - \text{arctan } x$ 
  apply (rule DERIV-neg-imp-decreasing-at-top [OF - *])
  apply (metis d-delta-arctan-upper-14 d-delta-arctan-upper-14-neg)
  done
then show ?thesis
  by auto
qed

```

2.2 Lower Bound 1

definition arctan-lower-11 :: real \Rightarrow real
where arctan-lower-11 $\equiv \lambda x. -(pi/2) - 3*x / (1 + 3*x^2)$

lemma arctan-lower-11:
assumes $x < 0$
shows $\text{arctan}(x) > \text{arctan-lower-11 } x$
using arctan-upper-14 [of $-x$] *assms*
by (auto simp: arctan-upper-14-def arctan-lower-11-def arctan-minus)

abbreviation arctan-lower-12 \equiv arctan-upper-13

lemma arctan-lower-12:
assumes $x \leq 0$
shows $\text{arctan}(x) \geq \text{arctan-lower-12 } x$
using arctan-upper-13 [of $-x$] *assms*
by (auto simp: arctan-upper-13-def arctan-minus)

abbreviation arctan-lower-13 \equiv arctan-upper-12

lemma arctan-lower-13:
assumes $x \geq 0$
shows $\text{arctan}(x) \geq \text{arctan-lower-13 } x$
using arctan-upper-12 [of $-x$] *assms*
by (auto simp: arctan-upper-12-def arctan-minus)

definition arctan-lower-14 :: real \Rightarrow real
where arctan-lower-14 $\equiv \lambda x. pi/2 - 1/x$

lemma arctan-lower-14:
assumes $x > 0$
shows $\text{arctan}(x) > \text{arctan-lower-14 } x$
using arctan-upper-11 [of $-x$] *assms*
by (auto simp: arctan-upper-11-def arctan-lower-14-def arctan-minus)

2.3 Upper Bound 3

```

definition arctan-upper-31 :: real  $\Rightarrow$  real
  where arctan-upper-31  $\equiv \lambda x. -(pi/2) - (64 + 735*x^2 + 945*x^4) / (15*x*(15 + 70*x^2 + 63*x^4))$ 

definition diff-delta-arctan-upper-31 :: real  $\Rightarrow$  real
  where diff-delta-arctan-upper-31  $\equiv \lambda x. 64 / (x^2 * (15 + 70*x^2 + 63*x^4))^2 * (1 + x^2)$ 

lemma d-delta-arctan-upper-31:
  assumes  $x \neq 0$ 
  shows  $((\lambda x. \text{arctan-upper-31 } x - \text{arctan } x) \text{ has-field-derivative } \text{diff-delta-arctan-upper-31}$ 
 $x)$  (at  $x$ )
  unfolding arctan-upper-31-def diff-delta-arctan-upper-31-def
  using assms
  apply (intro derivative-eq-intros)
  apply (rule refl | simp add: add-nonneg-eq-0-iff)+
  apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
  done

lemma d-delta-arctan-upper-31-pos:  $x \neq 0 \implies \text{diff-delta-arctan-upper-31 } x > 0$ 
unfolding diff-delta-arctan-upper-31-def
by (auto simp: divide-simps zero-less-mult-iff add-pos-pos add-nonneg-eq-0-iff)

lemma arctan-upper-31:
  assumes  $x < 0$ 
  shows  $\text{arctan}(x) < \text{arctan-upper-31 } x$ 
proof -
  have *:  $\bigwedge x:\text{real}. (15 + 70 * x^2 + 63 * x^4) > 0$ 
  by (sos ((R<1 + ((R<1 * ((R<7/8 * [19/7*x^2 + 1]^2) + ((R<4 * [x]^2) + (R<10/7 * [x^2]^2)))) + ((A<=0 * R<1) * (R<1/8 * [1]^2))))) + ((A<=0 * R<1) * (R<1/8 * [1]^2)))
  then have **:  $\bigwedge x:\text{real}. \neg (15 + 70 * x^2 + 63 * x^4) < 0$ 
  by (simp add: not-less)
  have  $((\lambda x:\text{real}. (64 + 735 * x^2 + 945 * x^4) / (15 * x * (15 + 70 * x^2 + 63 * x^4))) \longrightarrow 0)$  at-bot
  apply (rule tendsto-0-le [where  $f = \text{inverse}$  and  $K=2$ ])
  apply (metis at-bot-le-at-infinity tendsto-inverse-0 tendsto-mono)
  apply (simp add: eventually-at-bot-linorder)
  apply (rule-tac  $x=-1$  in exI)
  apply (auto simp: divide-simps abs-if zero-less-mult-iff **)
  done
  then have  $((\lambda x. \text{arctan-upper-31 } x - \text{arctan } x) \longrightarrow - (pi / 2) - 0 - (- (pi / 2)))$  at-bot
  unfolding arctan-upper-31-def
  apply (intro tendsto-intros tendsto-arctan-at-bot, auto)
  done
  then have *:  $((\lambda x. \text{arctan-upper-31 } x - \text{arctan } x) \longrightarrow 0)$  at-bot
  by simp

```

```

have 0 < arctan-upper-31 x - arctan x
  apply (rule DERIV-pos-imp-increasing-at-bot [OF - *])
  apply (metis assms d-delta-arctan-upper-31 d-delta-arctan-upper-31-pos not-le)
  done
then show ?thesis
  by auto
qed

definition arctan-upper-32 :: real ⇒ real
  where arctan-upper-32 ≡ λx. 7*(33*x^4 + 170*x^2 + 165)*x / (5*(5*x^6 +
105*x^4 + 315*x^2 + 231))

definition diff-delta-arctan-upper-32 :: real ⇒ real
  where diff-delta-arctan-upper-32 ≡ λx. -256*x^12 / ((5*x^6+105*x^4+315*x^2+231)^2*(1+x^2))

lemma d-delta-arctan-upper-32:
  ((λx. arctan-upper-32 x - arctan x) has-field-derivative diff-delta-arctan-upper-32
  x) (at x)
  unfolding arctan-upper-32-def diff-delta-arctan-upper-32-def
  apply (intro derivative-eq-intros | simp)+
  apply simp-all
  apply (auto simp: add-nonneg-eq-0-iff divide-simps, algebra)
  done

lemma arctan-upper-32:
  assumes x ≤ 0 shows arctan(x) ≤ arctan-upper-32 x
  apply (rule gen-upper-bound-decreasing [OF assms d-delta-arctan-upper-32])
  apply (auto simp: diff-delta-arctan-upper-32-def arctan-upper-32-def)
  done

definition arctan-upper-33 :: real ⇒ real
  where arctan-upper-33 ≡ λx. (64*x^4+735*x^2+945)*x / (15*(15*x^4+70*x^2+63))

definition diff-delta-arctan-upper-33 :: real ⇒ real
  where diff-delta-arctan-upper-33 ≡ λx. 64*x^10 / ((15*x^4+70*x^2+63)^2*(1+x^2))

lemma d-delta-arctan-upper-33:
  ((λx. arctan-upper-33 x - arctan x) has-field-derivative diff-delta-arctan-upper-33
  x) (at x)
  unfolding arctan-upper-33-def diff-delta-arctan-upper-33-def
  apply (intro derivative-eq-intros, simp-all)
  apply (auto simp: add-nonneg-eq-0-iff divide-simps, algebra)
  done

lemma arctan-upper-33:
  assumes x ≥ 0 shows arctan(x) ≤ arctan-upper-33 x
  apply (rule gen-upper-bound-increasing [OF assms d-delta-arctan-upper-33])
  apply (auto simp: diff-delta-arctan-upper-33-def arctan-upper-33-def)
  done

```

```

definition arctan-upper-34 :: real ⇒ real
  where arctan-upper-34 ≡
    λx. pi/2 - (33 + 170*x^2 + 165*x^4)*7*x / (5*(5 + 105*x^2 + 315*x^4
+ 231*x^6))

definition diff-delta-arctan-upper-34 :: real ⇒ real
  where diff-delta-arctan-upper-34 ≡ λx. -256 / ((5+105*x^2+315*x^4+231*x^6)^2*(1+x^2))

lemma d-delta-arctan-upper-34:
  ((λx. arctan-upper-34 x - arctan x) has-field-derivative diff-delta-arctan-upper-34
  x) (at x)
  unfolding arctan-upper-34-def diff-delta-arctan-upper-34-def
  apply (intro derivative-eq-intros | simp add: add-nonneg-eq-0-iff) +
  apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
  done

lemma d-delta-arctan-upper-34-pos: diff-delta-arctan-upper-34 x < 0
  unfolding diff-delta-arctan-upper-34-def
  apply (simp add: divide-simps add-nonneg-eq-0-iff zero-less-mult-iff)
  using power2-less-0 [of x]
  apply arith
  done

lemma arctan-upper-34:
  assumes x > 0
  shows arctan(x) < arctan-upper-34 x
  proof -
    have ((λx. arctan-upper-34 x - arctan x) ⟶ pi / 2 - 0 - pi / 2) at-top
    unfolding arctan-upper-34-def
    apply (intro tendsto-intros tendsto-arctan-at-top, auto)
    apply (rule tendsto-0-le [where f = inverse and K=1])
    apply (metis tendsto-inverse-0 at-top-le-at-infinity tendsto-mono)
    apply (simp add: eventually-at-top-linorder)
    apply (rule-tac x=1 in exI)
    apply (auto simp: divide-simps power-eq-if add-pos-pos algebra-simps)
    done
    then have *: ((λx. arctan-upper-34 x - arctan x) ⟶ 0) at-top
      by simp
    have 0 < arctan-upper-34 x - arctan x
      apply (rule DERIV-neg-imp-decreasing-at-top [OF - *])
      apply (metis d-delta-arctan-upper-34 d-delta-arctan-upper-34-pos)
      done
    then show ?thesis
      by auto
  qed

```

2.4 Lower Bound 3

definition *arctan-lower-31* :: *real* \Rightarrow *real*
where *arctan-lower-31* $\equiv \lambda x. -(pi/2) - (33 + 170*x^2 + 165*x^4)*7*x / (5*(5 + 105*x^2 + 315*x^4 + 231*x^6))$

lemma *arctan-lower-31*:
assumes $x < 0$
shows *arctan(x)* $>$ *arctan-lower-31 x*
using *arctan-upper-34* [of $-x$] *assms*
by (auto simp: *arctan-upper-34-def arctan-lower-31-def arctan-minus*)

abbreviation *arctan-lower-32* \equiv *arctan-upper-33*

lemma *arctan-lower-32*:
assumes $x \leq 0$
shows *arctan(x)* \geq *arctan-lower-32 x*
using *arctan-upper-33* [of $-x$] *assms*
by (auto simp: *arctan-upper-33-def arctan-minus*)

abbreviation *arctan-lower-33* \equiv *arctan-upper-32*

lemma *arctan-lower-33*:
assumes $x \geq 0$
shows *arctan(x)* \geq *arctan-lower-33 x*
using *arctan-upper-32* [of $-x$] *assms*
by (auto simp: *arctan-upper-32-def arctan-minus*)

definition *arctan-lower-34* :: *real* \Rightarrow *real*
where *arctan-lower-34* $\equiv \lambda x. pi/2 - (64 + 735*x^2 + 945*x^4) / (15*x*(15 + 70*x^2 + 63*x^4))$

lemma *arctan-lower-34*:
assumes $x > 0$
shows *arctan(x)* $>$ *arctan-lower-34 x*
using *arctan-upper-31* [of $-x$] *assms*
by (auto simp: *arctan-upper-31-def arctan-lower-34-def arctan-minus*)

2.5 Upper Bound 4

definition *arctan-upper-41* :: *real* \Rightarrow *real*
where *arctan-upper-41* \equiv

$$\lambda x. -(pi/2) - (256 + 5943*x^2 + 19250*x^4 + 15015*x^6) / (35*x*(35 + 315*x^2 + 693*x^4 + 429*x^6))$$

definition *diff-delta-arctan-upper-41* :: *real* \Rightarrow *real*
where *diff-delta-arctan-upper-41* $\equiv \lambda x. 256 / (x^2*(35+315*x^2+693*x^4+429*x^6)^2*(1+x^2))$

lemma *d-delta-arctan-upper-41*:

```

assumes  $x \neq 0$ 
shows  $((\lambda x. \text{arctan-upper-41 } x - \text{arctan } x) \text{ has-field-derivative } \text{diff-delta-arctan-upper-41}$   

 $x)$  (at  $x$ )
unfolding  $\text{arctan-upper-41-def } \text{diff-delta-arctan-upper-41-def}$ 
using assms
apply (intro derivative-eq-intros)
apply (rule refl | simp add: add-nonneg-eq-0-iff)+
apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
done

lemma  $d\text{-delta-arctan-upper-41-pos}: x \neq 0 \implies \text{diff-delta-arctan-upper-41 } x > 0$ 
unfolding  $\text{diff-delta-arctan-upper-41-def}$ 
by (auto simp: zero-less-mult-iff add-pos-pos add-nonneg-eq-0-iff)

lemma  $\text{arctan-upper-41}:$ 
assumes  $x < 0$ 
shows  $\text{arctan}(x) < \text{arctan-upper-41 } x$ 
proof -
have *:  $\bigwedge x:\text{real}. (35 + 315 * x^2 + 693 * x^4 + 429 * x^6) > 0$ 
by (sos (( $R < 1$ ) + (( $R < 13/8589934592 * [95/26*x^2 + 1]^2$ ) +  

( $(R < 38654705675/4294967296 * [170080704731/154618822700*x^3 + x]^2$ ) +  

( $(R < 14271/446676598784 * [x^2]^2$ ) + ( $(R < 3631584276674589067439/2656331147370089676800$   

*  $[x^3]^2$ )))) + (( $A \leq 0 * R < 1$ ) * ( $R < 245426703/8589934592 * [1]^2$ ))))))
then have **:  $\bigwedge x:\text{real}. x < 0 \implies \neg (35 + 315 * x^2 + 693 * x^4 + 429 * x^6) < 0$ 
by (simp add: not-less)
have  $((\lambda x:\text{real}. (256 + 5943 * x^2 + 19250 * x^4 + 15015 * x^6) /$ 
 $(35 * x * (35 + 315 * x^2 + 693 * x^4 + 429 * x^6))) \longrightarrow 0)$  at-bot
apply (rule tendsto-0-le [where  $f = \text{inverse}$  and  $K=2$ ])
apply (metis at-bot-le-at-infinity tendsto-inverse-0 tendsto-mono)
apply (simp add: eventually-at-bot-linorder)
apply (rule-tac  $x=-1$  in exI)
apply (auto simp: ** abs-if divide-simps zero-less-mult-iff)
done
then have  $((\lambda x. \text{arctan-upper-41 } x - \text{arctan } x) \longrightarrow -(pi/2) - 0 - (- (pi/2)))$  at-bot
unfolding  $\text{arctan-upper-41-def}$ 
apply (intro tendsto-intros tendsto-arctan-at-bot, auto)
done
then have *:  $((\lambda x. \text{arctan-upper-41 } x - \text{arctan } x) \longrightarrow 0)$  at-bot
by simp
have  $0 < \text{arctan-upper-41 } x - \text{arctan } x$ 
apply (rule DERIV-pos-imp-increasing-at-bot [OF -*])
apply (metis assms d-delta-arctan-upper-41 d-delta-arctan-upper-41-pos not-le)
done
then show ?thesis
by auto
qed

```

```

definition arctan-upper-42 :: real  $\Rightarrow$  real
where arctan-upper-42  $\equiv$ 

$$\lambda x. (15159*x^6 + 147455*x^4 + 345345*x^2 + 225225)*x / (35*(35*x^8 + 1260*x^6 + 6930*x^4 + 12012*x^2))$$


definition diff-delta-arctan-upper-42 :: real  $\Rightarrow$  real
where diff-delta-arctan-upper-42  $\equiv$ 

$$\lambda x. -16384*x^16 / ((35*x^8 + 1260*x^6 + 6930*x^4 + 12012*x^2 + 6435)^2 * (1+x^2))$$


lemma d-delta-arctan-upper-42:
 $((\lambda x. \text{arctan-upper-42 } x - \text{arctan } x) \text{ has-field-derivative } \text{diff-delta-arctan-upper-42}$ 
 $x) \text{ (at } x)$ 
unfolding arctan-upper-42-def diff-delta-arctan-upper-42-def
apply (intro derivative-eq-intros, simp-all)
apply (auto simp: divide-simps add-nonneg-eq-0-iff, algebra)
done

lemma arctan-upper-42:
assumes  $x \leq 0$  shows  $\text{arctan}(x) \leq \text{arctan-upper-42 } x$ 
apply (rule gen-upper-bound-decreasing [OF assms d-delta-arctan-upper-42])
apply (auto simp: diff-delta-arctan-upper-42-def arctan-upper-42-def)
done

definition arctan-upper-43 :: real  $\Rightarrow$  real
where arctan-upper-43  $\equiv$ 

$$\lambda x. (256*x^6 + 5943*x^4 + 19250*x^2 + 15015)*x / (35 * (35*x^6 + 315*x^4 + 693*x^2 + 429))$$


definition diff-delta-arctan-upper-43 :: real  $\Rightarrow$  real
where diff-delta-arctan-upper-43  $\equiv$   $\lambda x. 256*x^14 / ((35*x^6 + 315*x^4 + 693*x^2 + 429)^2 * (1+x^2))$ 

lemma d-delta-arctan-upper-43:
 $((\lambda x. \text{arctan-upper-43 } x - \text{arctan } x) \text{ has-field-derivative } \text{diff-delta-arctan-upper-43}$ 
 $x) \text{ (at } x)$ 
unfolding arctan-upper-43-def diff-delta-arctan-upper-43-def
apply (intro derivative-eq-intros, simp-all)
apply (auto simp: add-nonneg-eq-0-iff divide-simps, algebra)
done

lemma arctan-upper-43:
assumes  $x \geq 0$  shows  $\text{arctan}(x) \leq \text{arctan-upper-43 } x$ 
apply (rule gen-upper-bound-increasing [OF assms d-delta-arctan-upper-43])
apply (auto simp: diff-delta-arctan-upper-43-def arctan-upper-43-def)
done

definition arctan-upper-44 :: real  $\Rightarrow$  real
where arctan-upper-44  $\equiv$ 

$$\lambda x. \pi/2 - (15159 + 147455*x^2 + 345345*x^4 + 225225*x^6)*x / (35*(35 + 1260*x^2 + 6930*x^4 + 12012*x^6 + 6435*x^8))$$


```

```

definition diff-delta-arctan-upper-44 :: real ⇒ real
  where diff-delta-arctan-upper-44 ≡
    λx. −16384 / ((35 + 1260*x^2 + 6930*x^4 + 12012*x^6 + 6435*x^8)^2 * (1 + x^2))

lemma d-delta-arctan-upper-44:
  ((λx. arctan-upper-44 x − arctan x) has-field-derivative diff-delta-arctan-upper-44
  x) (at x)
  unfolding arctan-upper-44-def diff-delta-arctan-upper-44-def
  apply (intro derivative-eq-intros | simp add: add-nonneg-eq-0-iff) +
  apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
  done

lemma d-delta-arctan-upper-44-pos: diff-delta-arctan-upper-44 x < 0
  unfolding diff-delta-arctan-upper-44-def
  apply (auto simp: divide-simps add-nonneg-eq-0-iff zero-less-mult-iff)
  using power2-less-0 [of x]
  apply arith
  done

lemma arctan-upper-44:
  assumes x > 0
  shows arctan(x) < arctan-upper-44 x
proof −
  have ((λx. arctan-upper-44 x − arctan x) —→ pi / 2 − 0 − pi / 2) at-top
  unfolding arctan-upper-44-def
  apply (intro tendsto-intros tendsto-arctan-at-top, auto)
  apply (rule tendsto-0-le [where f = inverse and K=1])
  apply (metis tendsto-inverse-0 at-top-le-at-infinity tendsto-mono)
  apply (simp add: eventually-at-top-linorder)
  apply (rule-tac x=1 in ext)
  apply (auto simp: zero-le-mult-iff divide-simps not-le[symmetric] power-eq-if
  algebra-simps)
  done
  then have *: ((λx. arctan-upper-44 x − arctan x) —→ 0) at-top
  by simp
  have 0 < arctan-upper-44 x − arctan x
  apply (rule DERIV-neg-imp-decreasing-at-top [OF - *])
  apply (metis d-delta-arctan-upper-44 d-delta-arctan-upper-44-pos)
  done
  then show ?thesis
  by auto
qed

```

2.6 Lower Bound 4

```

definition arctan-lower-41 :: real ⇒ real
  where arctan-lower-41 ≡
    λx. −(pi/2) − (15159 + 147455*x^2 + 345345*x^4 + 225225*x^6)*x /
      (35*(35 + 1260*x^2 + 6930*x^4 + 12012*x^6 + 6435*x^8))

```

```

lemma arctan-lower-41:
  assumes x < 0
  shows arctan(x) > arctan-lower-41 x
  using arctan-upper-44 [of -x] assms
  by (auto simp: arctan-upper-44-def arctan-lower-41-def arctan-minus)

abbreviation arctan-lower-42 ≡ arctan-upper-43

lemma arctan-lower-42:
  assumes x ≤ 0
  shows arctan(x) ≥ arctan-lower-42 x
  using arctan-upper-43 [of -x] assms
  by (auto simp: arctan-upper-43-def arctan-minus)

abbreviation arctan-lower-43 ≡ arctan-upper-42

lemma arctan-lower-43:
  assumes x ≥ 0
  shows arctan(x) ≥ arctan-lower-43 x
  using arctan-upper-42 [of -x] assms
  by (auto simp: arctan-upper-42-def arctan-minus)

definition arctan-lower-44 :: real ⇒ real
  where arctan-lower-44 ≡
    λx. pi/2 - (256+5943*x^2+19250*x^4+15015*x^6) /
      (35*x*(35+315*x^2+693*x^4+429*x^6))

lemma arctan-lower-44:
  assumes x > 0
  shows arctan(x) > arctan-lower-44 x
  using arctan-upper-41 [of -x] assms
  by (auto simp: arctan-upper-41-def arctan-lower-44-def arctan-minus)

end

```

Chapter 3

Exp Upper and Lower Bounds

```
theory Exp-Bounds
imports Bounds-Lemmas
HOL-Library.Sum-of-Squares
Sturm-Sequences.Sturm
```

```
begin
```

Covers all bounds used in exp-upper.ax, exp-lower.ax and exp-extended.ax.

3.1 Taylor Series Bounds

exp-positive is the theorem $0 \leq \exp ?x$

exp-lower-taylor-1 is the theorem $1 + ?x \leq \exp ?x$

All even approximants are lower bounds.

```
lemma exp-lower-taylor-even:
fixes x::real
shows even n ==> ( $\sum m < n. (x^m) / (\text{fact } m)$ )  $\leq \exp x$ 
using Maclaurin-exp-le [of x n]
by (auto simp add: zero-le-even-power)
```

```
lemma exp-upper-taylor-even:
fixes x::real
assumes n: even n
and pos: ( $\sum m < n. ((-x)^m) / (\text{fact } m)$ )  $> 0$  (is ?sum > 0)
shows exp x  $\leq \text{inverse } ?sum$ 
using exp-lower-taylor-even [OF n, of -x]
by (metis exp-minus inverse-inverse-eq le-imp-inverse-le pos)
```

3 if the previous lemma is expressed in terms of $(2::'a) * m$.

```
lemma exp-lower-taylor-3:
```

```

fixes x::real
shows 1 + x + (1/2)*x2 + (1/6)*x3 + (1/24)*x4 + (1/120)*x5 ≤ exp
x
by (rule order-trans [OF - exp-lower-taylor-even [of 6]])
(auto simp: lessThan-nat-numeral fact-numeral)

lemma exp-lower-taylor-3-cubed:
fixes x::real
shows (1 + x/3 + (1/2)*(x/3)2 + (1/6)*(x/3)3 + (1/24)*(x/3)4 +
(1/120)*(x/3)5)3 ≤ exp x
proof -
  have (1 + x/3 + (1/2)*(x/3)2 + (1/6)*(x/3)3 + (1/24)*(x/3)4 +
(1/120)*(x/3)5)3
  ≤ exp (x/3)3
  by (metis power-mono-odd odd-numeral exp-lower-taylor-3)
  also have ... = exp x
  by (simp add: exp-of-nat-mult [symmetric])
  finally show ?thesis .
qed

lemma exp-lower-taylor-2:
fixes x::real
shows 1 + x + (1/2)*x2 + (1/6)*x3 ≤ exp x
proof -
  have even (4::nat) by simp
  then have (∑ m<4. xm / (fact m)) ≤ exp x
  by (rule exp-lower-taylor-even)
  then show ?thesis by (auto simp add: numeral-eq-Suc)
qed

lemma exp-upper-bound-case-3:
fixes x::real
assumes x ≤ 3.19
shows exp x ≤ 2304 / (-(x3) + 6*x2 - 24*x + 48)2
proof -
  have (1/48)*(-(x3) + 6*x2 - 24*x + 48) = (1 + (-x/2)) + (1/2)*(-x/2)2
  + (1/6)*(-x/2)3
  by (simp add: field-simps power2-eq-square power3-eq-cube)
  also have ... ≤ exp (-x/2)
  by (rule exp-lower-taylor-2)
  finally have 1: (1/48)*(-(x3) + 6*x2 - 24*x + 48) ≤ exp (-x/2) .
  have (-(x3) + 6*x2 - 24*x + 48)2 / 2304 = ((1/48)*(-(x3) + 6*x2
  - 24*x + 48))2
  by (simp add: field-simps power2-eq-square power3-eq-cube)
  also have ... ≤ (exp (-x/2))2
  apply (rule power-mono [OF 1])
  apply (simp add: algebra-simps)
  using assms
  apply (sos ((R<1 + ((R<1 * ((R<1323/13 * [~15/49*x + 1])2) + (R<1/637

```

```

* [x] ^2))) + (((A < 0 * R < 1) * (R < 50 / 13 * [1] ^2)) + ((A <= 0 * R < 1) * ((R < 56 / 13
* [~5 / 56 * x + 1] ^2) + (R < 199 / 728 * [x] ^2)))))))
done
also have ... = inverse (exp x)
by (metis exp-minus mult-exp-exp power2-eq-square field-sum-of-halves)
finally have 2: (- (x ^3) + 6 * x ^2 - 24 * x + 48) ^2 / 2304 ≤ inverse (exp x) .
have 6 * x ^2 - x ^3 - 24 * x + 48 ≠ 0 using assms
by (sos ((R < 1 + ([~400 / 13] * A = 0) + ((R < 1 * ((R < 1323 / 13 * [~15 / 49 * x
+ 1] ^2) + (R < 1 / 637 * [x] ^2))) + ((A <= 0 * R < 1) * ((R < 56 / 13 * [~5 / 56 * x
+ 1] ^2) + (R < 199 / 728 * [x] ^2)))))))
then show ?thesis
using Fields.linordered-field-class.le-imp-inverse-le [OF 2]
by simp
qed

```

```

lemma exp-upper-bound-case-5:
fixes x::real
assumes x ≤ 6.36
shows exp x ≤ 21743271936 / (- (x ^3) + 12 * x ^2 - 96 * x + 384) ^4
proof –
have (1 / 384) * (- (x ^3) + 12 * x ^2 - 96 * x + 384) = (1 + (-x / 4) + (1 / 2) * (-x / 4) ^2
+ (1 / 6) * (-x / 4) ^3)
by (simp add: field-simps power2-eq-square power3-eq-cube)
also have ... ≤ exp (-x / 4)
by (rule exp-lower-taylor-2)
finally have 1: (1 / 384) * (- (x ^3) + 12 * x ^2 - 96 * x + 384) ≤ exp (-x / 4) .
have (- (x ^3) + 12 * x ^2 - 96 * x + 384) ^4 / 21743271936 = ((1 / 384) * (- (x ^3)
+ 12 * x ^2 - 96 * x + 384)) ^4
by (simp add: divide-simps)
also have ... ≤ (exp (-x / 4)) ^4
apply (rule power-mono [OF 1])
apply (simp add: algebra-simps)
using assms
apply (sos ((R < 1 + ((R < 1 * ((R < 1777 / 32 * [~539 / 3554 * x + 1] ^2) +
(R < 907 / 227456 * [x] ^2))) + (((A < 0 * R < 1) * (R < 25 / 1024 * [1] ^2)) + ((A <= 0
* R < 1) * ((R < 49 / 32 * [~2 / 49 * x + 1] ^2) + (R < 45 / 1568 * [x] ^2)))))))
done
also have ... = inverse (exp x)
by (simp add: exp-of-nat-mult [symmetric] exp-minus [symmetric])
finally have 2: (- (x ^3) + 12 * x ^2 - 96 * x + 384) ^4 / 21743271936 ≤ inverse
(exp x) .
have 12 * x ^2 - x ^3 - 96 * x + 384 ≠ 0 using assms
by (sos ((R < 1 + ([~25 / 32] * A = 0) + ((R < 1 * ((R < 1777 / 32 * [~539 / 3554 * x
+ 1] ^2) + (R < 907 / 227456 * [x] ^2))) + ((A <= 0 * R < 1) * ((R < 49 / 32 * [~2 / 49 * x
+ 1] ^2) + (R < 45 / 1568 * [x] ^2)))))))
then show ?thesis
using Fields.linordered-field-class.le-imp-inverse-le [OF 2]
by simp
qed

```

3.2 Continued Fraction Bound 2

```

definition exp-cf2 :: real  $\Rightarrow$  real
  where exp-cf2  $\equiv \lambda x. (x^2 + 6*x + 12) / (x^2 - 6*x + 12)$ 

lemma denom-cf2-pos: fixes x::real shows  $x^2 - 6*x + 12 > 0$ 
  by (sos ((R<1 + ((R<1 * ((R<5 * [~3/10*x + 1]^2) + (R<1/20 * [x]^2))) + ((A<=0 * R<1) * (R<1/2 * [1]^2)))))

lemma numer-cf2-pos: fixes x::real shows  $x^2 + 6*x + 12 > 0$ 
  by (sos ((R<1 + ((R<1 * ((R<5 * [3/10*x + 1]^2) + (R<1/20 * [x]^2))) + ((A<=0 * R<1) * (R<1/2 * [1]^2)))))

lemma exp-cf2-pos: exp-cf2 x > 0
  unfolding exp-cf2-def
  by (auto simp add: divide-simps denom-cf2-pos numer-cf2-pos)

definition diff-delta-lnexp-cf2 :: real  $\Rightarrow$  real
  where diff-delta-lnexp-cf2  $\equiv \lambda x. - (x^4) / ((x^2 - 6*x + 12) * (x^2 + 6*x + 12))$ 

lemma d-delta-lnexp-cf2-nonpos: diff-delta-lnexp-cf2 x  $\leq 0$ 
  unfolding diff-delta-lnexp-cf2-def
  by (sos (((R<1 + ((R<1 * ((R<5/4 * [~3/40*x^2 + 1]^2) + (R<11/1280 * [x^2]^2)))) + ((A<1 * R<1) * (R<1/64 * [1]^2)))) & ((R<1 + ((R<1 * ((R<5/4 * [~3/40*x^2 + 1]^2) + (R<11/1280 * [x^2]^2))) + ((A<1 * R<1) * (R<1/64 * [1]^2)))))

lemma d-delta-lnexp-cf2:
   $((\lambda x. \ln(\exp-cf2 x) - x) \text{ has-field-derivative } \text{diff-delta-lnexp-cf2 } x) \text{ (at } x)$ 
  unfolding exp-cf2-def diff-delta-lnexp-cf2-def
  apply (intro derivative-eq-intros | simp)+
  apply (metis exp-cf2-def exp-cf2-pos)
  apply (simp-all add:)
  using denom-cf2-pos [of x] numer-cf2-pos [of x]
  apply (auto simp: divide-simps)
  apply algebra
  done

  Upper bound for non-positive x

lemma ln-exp-cf2-upper-bound-neg:
  assumes x  $\leq 0$ 
  shows x  $\leq \ln(\exp-cf2 x)$ 
  by (rule gen-upper-bound-decreasing [OF assms d-delta-lnexp-cf2 d-delta-lnexp-cf2-nonpos])
    (simp add: exp-cf2-def)

theorem exp-cf2-upper-bound-neg:  $x \leq 0 \implies \exp(x) \leq \exp-cf2 x$ 
  by (metis ln-exp-cf2-upper-bound-neg exp-cf2-pos exp-le-cancel-iff exp-ln-iff)

```

Lower bound for non-negative x

```
lemma ln-exp-cf2-lower-bound-pos:
  assumes 0 ≤ x
  shows ln (exp-cf2 x) ≤ x
  by (rule gen-lower-bound-increasing [OF assms d-delta-lnexp-cf2 d-delta-lnexp-cf2-nonpos])
    (simp add: exp-cf2-def)

theorem exp-cf2-lower-bound-pos: 0 ≤ x ⇒ exp-cf2 x ≤ exp x
  by (metis exp-cf2-pos exp-le-cancel-iff exp-ln ln-exp-cf2-lower-bound-pos)
```

3.3 Continued Fraction Bound 3

This bound crosses the X-axis twice, causing complications.

```
definition numer-cf3 :: real ⇒ real
  where numer-cf3 ≡ λx. x^3 + 12*x^2 + 60*x + 120
```

```
definition exp-cf3 :: real ⇒ real
  where exp-cf3 ≡ λx. numer-cf3 x / numer-cf3 (-x)
```

```
lemma numer-cf3-pos: -4.64 ≤ x ⇒ numer-cf3 x > 0
  unfolding numer-cf3-def
  by sturm
```

```
lemma exp-cf3-pos: numer-cf3 x > 0 ⇒ numer-cf3 (-x) > 0 ⇒ exp-cf3 x > 0
  by (simp add: exp-cf3-def)
```

```
definition diff-delta-lnexp-cf3 :: real ⇒ real
  where diff-delta-lnexp-cf3 ≡ λx. (x^6) / (numer-cf3 (-x) * numer-cf3 x)
```

```
lemma d-delta-lnexp-cf3-nonneg: numer-cf3 x > 0 ⇒ numer-cf3 (-x) > 0 ⇒
diff-delta-lnexp-cf3 x ≥ 0
  unfolding diff-delta-lnexp-cf3-def
  by (auto simp: mult-less-0-iff intro: divide-nonneg-neg)
```

```
lemma d-delta-lnexp-cf3:
  assumes numer-cf3 x > 0 numer-cf3 (-x) > 0
  shows ((λx. ln (exp-cf3 x) - x) has-field-derivative diff-delta-lnexp-cf3 x) (at x)
  unfolding exp-cf3-def numer-cf3-def diff-delta-lnexp-cf3-def
  apply (intro derivative-eq-intros | simp)+
  using assms numer-cf3-pos [of x] numer-cf3-pos [of -x]
  apply (auto simp: numer-cf3-def)
  apply (auto simp add: divide-simps add-nonneg-eq-0-iff)
  apply algebra
  done
```

```
lemma numer-cf3-mono: y ≤ x ⇒ numer-cf3 y ≤ numer-cf3 x
  unfolding numer-cf3-def
```

by (sos (((A<0 * R<1) + ((A<=0 * R<1) * ((R<60 * [1/10*x + 1/10*y + 1]^2) + ((R<2/5 * [x + ~1/4*y]^2) + (R<3/8 * [y]^2)))))))

Upper bound for non-negative x

lemma ln-exp-cf3-upper-bound-nonneg:

assumes x0: $0 \leq x$ **and** xless: numer-cf3 ($-x$) > 0

shows $x \leq \ln(\exp\text{-}cf3 x)$

proof –

have ncf3: $\forall y. 0 \leq y \implies y \leq x \implies \text{numer-cf3 } (-y) > 0$

by (metis neg-le-iff-le numer-cf3-mono order.strict-trans2 xless)

show ?thesis

apply (rule gen-upper-bound-increasing [OF x0 d-delta-lnexp-cf3 d-delta-lnexp-cf3-nonneg])

apply (auto simp add: ncf3 assms numer-cf3-pos)

apply (simp add: exp-cf3-def numer-cf3-def)

done

qed

theorem exp-cf3-upper-bound-pos: $0 \leq x \implies \text{numer-cf3 } (-x) > 0 \implies \exp x \leq \exp\text{-}cf3 x$

using ln-exp-cf3-upper-bound-nonneg [of x] exp-cf3-pos [of x] numer-cf3-pos [of x]

by auto (metis exp-le-cancel-iff exp-ln-iff)

corollary $0 \leq x \implies x \leq 4.64 \implies \exp x \leq \exp\text{-}cf3 x$

by (metis numer-cf3-pos neg-le-iff-le exp-cf3-upper-bound-pos)

Lower bound for negative x, provided $0 < \exp\text{-}cf3 x$

lemma ln-exp-cf3-lower-bound-neg:

assumes x0: $x \leq 0$ **and** xgtr: numer-cf3 x > 0

shows $\ln(\exp\text{-}cf3 x) \leq x$

proof –

have ncf3: $\forall y. x \leq y \implies y \leq 0 \implies \text{numer-cf3 } y > 0$

by (metis dual-order.strict-trans1 numer-cf3-mono xgtr)

show ?thesis

apply (rule gen-lower-bound-decreasing [OF x0 d-delta-lnexp-cf3 d-delta-lnexp-cf3-nonneg])

apply (auto simp add: ncf3 assms numer-cf3-pos)

apply (simp add: exp-cf3-def numer-cf3-def)

done

qed

theorem exp-cf3-lower-bound-pos:

assumes $x \leq 0$ **shows** $\exp\text{-}cf3 x \leq \exp x$

proof (cases numer-cf3 x > 0)

case True

then have $\exp\text{-}cf3 x > 0$

using assms numer-cf3-pos [of $-x$]

by (auto simp: exp-cf3-pos)

then show ?thesis

```

using ln-exp-cf3-lower-bound-neg [of x] assms True
  by auto (metis exp-le-cancel-iff exp-ln-iff)
next
case False
then have exp-cf3 x ≤ 0
  using assms numer-cf3-pos [of -x]
  unfolding exp-cf3-def
    by (simp add: divide-nonpos-pos)
then show ?thesis
  by (metis exp-ge-zero order.trans)
qed

```

3.4 Continued Fraction Bound 4

Here we have the extended exp continued fraction bounds

```

definition numer-cf4 :: real ⇒ real
  where numer-cf4 ≡ λx. x^4 + 20*x^3 + 180*x^2 + 840*x + 1680

definition exp-cf4 :: real ⇒ real
  where exp-cf4 ≡ λx. numer-cf4 x / numer-cf4 (-x)

lemma numer-cf4-pos: fixes x::real shows numer-cf4 x > 0
  unfolding numer-cf4-def
  by (sos ((R<1 + ((R<1 * ((R<4469/256 * [1135/71504*x^2 + 4725/17876*x +
  1]^2) + ((R<3728645/18305024 * [536265/2982916*x^2 + x]^2) + (R<106265/24436047872
  * [x^2]^2)))) + ((A<=0 * R<1) * (R<45/4096 * [1]^2)))))

lemma exp-cf4-pos: exp-cf4 x > 0
  unfolding exp-cf4-def
  by (auto simp add: divide-simps numer-cf4-pos)

definition diff-delta-lnexp-cf4 :: real ⇒ real
  where diff-delta-lnexp-cf4 ≡ λx. - (x^8) / (numer-cf4 (-x) * numer-cf4 x)

lemma d-delta-lnexp-cf4-nonpos: diff-delta-lnexp-cf4 x ≤ 0
  unfolding diff-delta-lnexp-cf4-def
  using numer-cf4-pos [of x] numer-cf4-pos [of -x]
  by (simp add: zero-le-divide-iff zero-le-mult-iff)

lemma d-delta-lnexp-cf4:
  ((λx. ln (exp-cf4 x) - x) has-field-derivative diff-delta-lnexp-cf4 x) (at x)
  unfolding exp-cf4-def numer-cf4-def diff-delta-lnexp-cf4-def
  apply (intro derivative-eq-intros | simp)+
  using exp-cf4-pos
  apply (simp add: exp-cf4-def numer-cf4-def)
  apply (simp-all add: )
  using numer-cf4-pos [of x] numer-cf4-pos [of -x]
  apply (auto simp: divide-simps numer-cf4-def)

```

```

apply algebra
done

    Upper bound for non-positive x

lemma ln-exp-cf4-upper-bound-neg:
  assumes  $x \leq 0$ 
  shows  $x \leq \ln(\exp-cf4 x)$ 
  by (rule gen-upper-bound-decreasing [OF assms d-delta-lnexp-cf4 d-delta-lnexp-cf4-nonpos])
    (simp add: exp-cf4-def numer-cf4-def)

theorem exp-cf4-upper-bound-neg:  $x \leq 0 \implies \exp(x) \leq \exp-cf4 x$ 
  by (metis ln-exp-cf4-upper-bound-neg exp-cf4-pos exp-le-cancel-iff exp-ln-iff)

    Lower bound for non-negative x

lemma ln-exp-cf4-lower-bound-pos:
  assumes  $0 \leq x$ 
  shows  $\ln(\exp-cf4 x) \leq x$ 
  by (rule gen-lower-bound-increasing [OF assms d-delta-lnexp-cf4 d-delta-lnexp-cf4-nonpos])
    (simp add: exp-cf4-def numer-cf4-def)

theorem exp-cf4-lower-bound-pos:  $0 \leq x \implies \exp-cf4 x \leq \exp x$ 
  by (metis exp-cf4-pos exp-le-cancel-iff exp-ln ln-exp-cf4-lower-bound-pos)

```

3.5 Continued Fraction Bound 5

This bound crosses the X-axis twice, causing complications.

```

definition numer-cf5 :: real ⇒ real
  where numer-cf5 ≡  $\lambda x. x^5 + 30*x^4 + 420*x^3 + 3360*x^2 + 15120*x + 30240$ 

definition exp-cf5 :: real ⇒ real
  where exp-cf5 ≡  $\lambda x. \text{numer-cf5 } x / \text{numer-cf5 } (-x)$ 

lemma numer-cf5-pos:  $-7.293 \leq x \implies \text{numer-cf5 } x > 0$ 
  unfolding numer-cf5-def
  by sturm

lemma exp-cf5-pos:  $\text{numer-cf5 } x > 0 \implies \text{numer-cf5 } (-x) > 0 \implies \exp-cf5 x > 0$ 
  unfolding exp-cf5-def numer-cf5-def
  by (simp add: divide-neg-neg)

definition diff-delta-lnexp-cf5 :: real ⇒ real
  where diff-delta-lnexp-cf5 ≡  $\lambda x. (x^{10}) / (\text{numer-cf5 } (-x) * \text{numer-cf5 } x)$ 

lemma d-delta-lnexp-cf5-nonneg:  $\text{numer-cf5 } x > 0 \implies \text{numer-cf5 } (-x) > 0 \implies$ 
 $\text{diff-delta-lnexp-cf5 } x \geq 0$ 
  unfolding diff-delta-lnexp-cf5-def
  by (auto simp add: mult-less-0-iff intro: divide-nonneg-neg )

```

```

lemma d-delta-lnexp-cf5:
  assumes numer-cf5 x > 0 numer-cf5 (-x) > 0
  shows (( $\lambda x$ . ln (exp-cf5 x) - x) has-field-derivative diff-delta-lnexp-cf5 x) (at x)
  unfolding exp-cf5-def numer-cf5-def diff-delta-lnexp-cf5-def
  apply (intro derivative-eq-intros | simp)+
  using assms numer-cf5-pos [of x] numer-cf5-pos [of -x]
  apply (auto simp: numer-cf5-def)
  apply (auto simp add: divide-simps add-nonneg-eq-0-iff)
  apply algebra
  done

```

3.5.1 Proving monotonicity via a non-negative derivative

```

definition numer-cf5-deriv :: real  $\Rightarrow$  real
  where numer-cf5-deriv  $\equiv$   $\lambda x$ .  $5*x^4 + 120*x^3 + 1260*x^2 + 6720*x + 15120$ 

```

```

lemma numer-cf5-deriv:
  shows (numer-cf5 has-field-derivative numer-cf5-deriv x) (at x)
  unfolding numer-cf5-def numer-cf5-deriv-def
  by (intro derivative-eq-intros | simp)+

lemma numer-cf5-deriv-pos: numer-cf5-deriv x  $\geq$  0
  unfolding numer-cf5-deriv-def
  by (sos ((R<1 + ((R<1 * ((R<185533/8192 * [73459/5937056*x^2 + 43050/185533*x + 1]^2) + ((R<4641265253/24318181376 * [700850925/4641265253*x^2 + x]^2) + (R<38142496079/38933754831437824 * [x^2]^2)))) + ((A<0 * R<1) * (R<205/131072 * [1]^2)))))

lemma numer-cf5-mono: y  $\leq$  x  $\implies$  numer-cf5 y  $\leq$  numer-cf5 x
  by (auto intro: DERIV-nonneg-imp-nondecreasing numer-cf5-deriv numer-cf5-deriv-pos)

```

3.5.2 Results

Upper bound for non-negative x

```

lemma ln-exp-cf5-upper-bound-nonneg:
  assumes x0: 0  $\leq$  x and xless: numer-cf5 (-x) > 0
  shows x  $\leq$  ln (exp-cf5 x)
  proof -
    have ncf5:  $\bigwedge y$ . 0  $\leq$  y  $\implies$  y  $\leq$  x  $\implies$  numer-cf5 (-y) > 0
      by (metis neg-le-iff-le numer-cf5-mono order.strict-trans2 xless)
    show ?thesis
      apply (rule gen-upper-bound-increasing [OF x0 d-delta-lnexp-cf5 d-delta-lnexp-cf5-nonneg])
      apply (auto simp add: ncf5 assms numer-cf5-pos)
      apply (simp add: exp-cf5-def numer-cf5-def)
      done
  qed

```

theorem *exp-cf5-upper-bound-pos*: $0 \leq x \implies \text{numer-cf5 } (-x) > 0 \implies \exp x \leq \exp\text{-cf5 } x$

using *ln-exp-cf5-upper-bound-nonneg* [of x] *exp-cf5-pos* [of x] *numer-cf5-pos* [of x]
by *auto* (*metis exp-le-cancel-iff exp-ln-iff*)

corollary $0 \leq x \implies x \leq 7.293 \implies \exp x \leq \exp\text{-cf5 } x$

by (*metis neg-le-iff-le numer-cf5-pos exp-cf5-upper-bound-pos*)

Lower bound for negative x , provided $0 < \exp\text{-cf5 } x]$

lemma *ln-exp-cf5-lower-bound-neg*:

assumes $x0: x \leq 0$ **and** $xgtr: \text{numer-cf5 } x > 0$
shows $\ln(\exp\text{-cf5 } x) \leq x$

proof –

have $ncf5: \bigwedge y. x \leq y \implies y \leq 0 \implies \text{numer-cf5 } y > 0$
by (*metis dual-order.strict-trans1 numer-cf5-mono xgtr*)
show ?thesis
apply (rule *gen-lower-bound-decreasing* [OF $x0 d\text{-delta-}lnexp\text{-cf5 } d\text{-delta-}lnexp\text{-cf5-nonneg}$])
apply (*auto simp add: ncf5 assms numer-cf5-pos*)
apply (*simp add: exp-cf5-def numer-cf5-def*)
done

qed

theorem *exp-cf5-lower-bound-pos*:

assumes $x \leq 0$ **shows** $\exp\text{-cf5 } x \leq \exp x$

proof (cases $\text{numer-cf5 } x > 0$)

case *True*
then have $\exp\text{-cf5 } x > 0$
using *assms numer-cf5-pos* [of $-x$]
by (*auto simp: exp-cf5-pos*)
then show ?thesis
using *ln-exp-cf5-lower-bound-neg* [of x] *assms True*
by *auto* (*metis exp-le-cancel-iff exp-ln-iff*)

next
case *False*
then have $\exp\text{-cf5 } x \leq 0$
using *assms numer-cf5-pos* [of $-x$]
unfolding *exp-cf5-def numer-cf5-def*
by (*simp add: divide-nonpos-pos*)
then show ?thesis
by (*metis exp-ge-zero order.trans*)

qed

3.6 Continued Fraction Bound 6

definition *numer-cf6* :: *real* \Rightarrow *real*
where $\text{numer-cf6} \equiv \lambda x. x^6 + 42*x^5 + 840*x^4 + 10080*x^3 + 75600*x^2 + 332640*x + 665280$

```

definition exp-cf6 :: real  $\Rightarrow$  real
  where exp-cf6  $\equiv \lambda x. \text{numer-cf6 } x / \text{numer-cf6 } (-x)$ 

lemma numer-cf6-pos: fixes x::real shows numer-cf6 x  $> 0$ 
  unfolding numer-cf6-def
  by sturm

lemma exp-cf6-pos: exp-cf6 x  $> 0$ 
  unfolding exp-cf6-def
  by (auto simp add: divide-simps numer-cf6-pos)

definition diff-delta-lnexp-cf6 :: real  $\Rightarrow$  real
  where diff-delta-lnexp-cf6  $\equiv \lambda x. - (x^{12}) / (\text{numer-cf6 } (-x) * \text{numer-cf6 } x)$ 

lemma d-delta-lnexp-cf6-nonpos: diff-delta-lnexp-cf6 x  $\leq 0$ 
  unfolding diff-delta-lnexp-cf6-def
  using numer-cf6-pos [of x] numer-cf6-pos [of -x]
  by (simp add: zero-le-divide-iff zero-le-mult-iff)

lemma d-delta-lnexp-cf6:
   $((\lambda x. \ln(\text{exp-cf6 } x) - x) \text{ has-field-derivative } \text{diff-delta-lnexp-cf6 } x) \text{ (at } x)$ 
  unfolding exp-cf6-def diff-delta-lnexp-cf6-def numer-cf6-def
  apply (intro derivative-eq-intros | simp)+
  using exp-cf6-pos
  apply (simp add: exp-cf6-def numer-cf6-def)
  apply (simp-all add: )
  using numer-cf6-pos [of x] numer-cf6-pos [of -x]
  apply (auto simp: divide-simps numer-cf6-def)
  apply algebra
  done

  Upper bound for non-positive x

lemma ln-exp-cf6-upper-bound-neg:
  assumes x  $\leq 0$ 
  shows x  $\leq \ln(\text{exp-cf6 } x)$ 
  by (rule gen-upper-bound-decreasing [OF assms d-delta-lnexp-cf6 d-delta-lnexp-cf6-nonpos])
    (simp add: exp-cf6-def numer-cf6-def)

theorem exp-cf6-upper-bound-neg:  $x \leq 0 \implies \exp(x) \leq \text{exp-cf6 } x$ 
  by (metis ln-exp-cf6-upper-bound-neg exp-cf6-pos exp-le-cancel-iff exp-ln-iff)

  Lower bound for non-negative x

lemma ln-exp-cf6-lower-bound-pos:
  assumes 0  $\leq x$ 
  shows  $\ln(\text{exp-cf6 } x) \leq x$ 
  by (rule gen-lower-bound-increasing [OF assms d-delta-lnexp-cf6 d-delta-lnexp-cf6-nonpos])
    (simp add: exp-cf6-def numer-cf6-def)

theorem exp-cf6-lower-bound-pos:  $0 \leq x \implies \exp(\text{exp-cf6 } x) \leq \exp x$ 
  by (metis exp-cf6-pos exp-le-cancel-iff exp-ln ln-exp-cf6-lower-bound-pos)

```

3.7 Continued Fraction Bound 7

This bound crosses the X-axis twice, causing complications.

```

definition numer-cf7 :: real  $\Rightarrow$  real
  where numer-cf7  $\equiv \lambda x. x^7 + 56*x^6 + 1512*x^5 + 25200*x^4 + 277200*x^3$ 
   $+ 1995840*x^2 + 8648640*x + 17297280$ 

definition exp-cf7 :: real  $\Rightarrow$  real
  where exp-cf7  $\equiv \lambda x. \text{numer-cf7 } x / \text{numer-cf7 } (-x)$ 

lemma numer-cf7-pos:  $-9.943 \leq x \implies \text{numer-cf7 } x > 0$ 
  unfolding numer-cf7-def
  by sturm

lemma exp-cf7-pos:  $\text{numer-cf7 } x > 0 \implies \text{numer-cf7 } (-x) > 0 \implies \text{exp-cf7 } x > 0$ 
  by (simp add: exp-cf7-def)

definition diff-delta-lnexp-cf7 :: real  $\Rightarrow$  real
  where diff-delta-lnexp-cf7  $\equiv \lambda x. (x^{14}) / (\text{numer-cf7 } (-x) * \text{numer-cf7 } x)$ 

lemma d-delta-lnexp-cf7-nonneg:  $\text{numer-cf7 } x > 0 \implies \text{numer-cf7 } (-x) > 0 \implies$ 
   $\text{diff-delta-lnexp-cf7 } x \geq 0$ 
  unfolding diff-delta-lnexp-cf7-def
  by (auto simp: mult-less-0-iff intro: divide-nonneg-neg)

lemma d-delta-lnexp-cf7:
  assumes numer-cf7 x > 0 numer-cf7 (-x) > 0
  shows (( $\lambda x. \ln(\text{exp-cf7 } x) - x$ ) has-field-derivative diff-delta-lnexp-cf7 x) (at x)
  unfolding exp-cf7-def numer-cf7-def diff-delta-lnexp-cf7-def
  apply (intro derivative-eq-intros | simp)+
  using assms numer-cf7-pos [of x] numer-cf7-pos [of -x]
  apply (auto simp: numer-cf7-def)
  apply (auto simp: divide-simps add-nonneg-eq-0-iff)
  apply algebra
  done

```

3.7.1 Proving monotonicity via a non-negative derivative

```

definition numer-cf7-deriv :: real  $\Rightarrow$  real
  where numer-cf7-deriv  $\equiv \lambda x. 7*x^6 + 336*x^5 + 7560*x^4 + 100800*x^3 +$ 
   $831600*x^2 + 3991680*x + 8648640$ 

lemma numer-cf7-deriv:
  shows (numer-cf7 has-field-derivative numer-cf7-deriv x) (at x)
  unfolding numer-cf7-def numer-cf7-deriv-def
  by (intro derivative-eq-intros | simp)+

lemma numer-cf7-deriv-pos:  $\text{numer-cf7-deriv } x \geq 0$ 
  unfolding numer-cf7-deriv-def

```

apply (*rule order.strict-implies-order*) — FIXME should not be necessary
by *sturm*

lemma *numer-cf7-mono*: $y \leq x \implies \text{numer-cf7 } y \leq \text{numer-cf7 } x$
by (*auto intro: DERIV-nonneg-imp-nondecreasing numer-cf7-deriv numer-cf7-deriv-pos*)

3.7.2 Results

Upper bound for non-negative x

lemma *ln-exp-cf7-upper-bound-nonneg*:
assumes *x0: 0 ≤ x and xless: numer-cf7 (-x) > 0*
shows *x ≤ ln (exp-cf7 x)*
proof –
have *ncf7: ∀y. 0 ≤ y ⇒ y ≤ x ⇒ numer-cf7 (-y) > 0*
 by (*metis neg-le-iff-le numer-cf7-mono order.strict-trans2 xless*)
show ?thesis
 apply (*rule gen-upper-bound-increasing [OF x0 d-delta-lnexp-cf7 d-delta-lnexp-cf7-nonneg]*)
 apply (*auto simp add: ncf7 assms numer-cf7-pos*)
 apply (*simp add: exp-cf7-def numer-cf7-def*)
 done
qed

theorem *exp-cf7-upper-bound-pos*: $0 \leq x \implies \text{numer-cf7 } (-x) > 0 \implies \exp x \leq \exp-cf7 x$
using *ln-exp-cf7-upper-bound-nonneg [of x] exp-cf7-pos [of x] numer-cf7-pos [of x]*
by *auto (metis exp-le-cancel-iff exp-ln-iff)*

corollary $0 \leq x \implies x \leq 9.943 \implies \exp x \leq \exp-cf7 x$
by (*metis neg-le-iff-le numer-cf7-pos exp-cf7-upper-bound-pos*)

Lower bound for negative x, provided $0 < \exp-cf7 x]$

lemma *ln-exp-cf7-lower-bound-neg*:
assumes *x0: x ≤ 0 and xgtr: numer-cf7 x > 0*
shows *ln (exp-cf7 x) ≤ x*
proof –
have *ncf7: ∀y. x ≤ y ⇒ y ≤ 0 ⇒ numer-cf7 y > 0*
 by (*metis dual-order.strict-trans1 numer-cf7-mono xgtr*)
show ?thesis
 apply (*rule gen-lower-bound-decreasing [OF x0 d-delta-lnexp-cf7 d-delta-lnexp-cf7-nonneg]*)
 apply (*auto simp add: ncf7 assms numer-cf7-pos*)
 apply (*simp add: exp-cf7-def numer-cf7-def*)
 done
qed

theorem *exp-cf7-lower-bound-pos*:
assumes *x ≤ 0 shows exp-cf7 x ≤ exp x*
proof (*cases numer-cf7 x > 0*)
case *True*

```

then have exp-cf7  $x > 0$ 
  using assms numer-cf7-pos [of  $-x$ ]
  by (auto simp: exp-cf7-pos)
then show ?thesis
  using ln-exp-cf7-lower-bound-neg [of  $x$ ] assms True
  by auto (metis exp-le-cancel-iff exp-ln-iff)
next
  case False
  then have exp-cf7  $x \leq 0$ 
    using assms numer-cf7-pos [of  $-x$ ]
    unfolding exp-cf7-def
    by (simp add: divide-nonpos-pos)
  then show ?thesis
    by (metis exp-ge-zero order.trans)
qed

end

```

Chapter 4

Log Upper and Lower Bounds

```
theory Log-CF-Bounds
imports Bounds-Lemmas

begin

theorem ln-upper-1: 0 < x ==> ln(x::real) ≤ x - 1
by (rule ln-le-minus-one)

definition ln-lower-1 :: real ⇒ real
where ln-lower-1 ≡ λx. 1 - (inverse x)

corollary ln-lower-1: 0 < x ==> ln-lower-1 x ≤ ln x
unfolding ln-lower-1-def
by (metis ln-inverse ln-le-minus-one positive-imp-inverse-positive minus-diff-eq
minus-le-iff)

theorem ln-lower-1-eq: 0 < x ==> ln-lower-1 x = (x - 1)/x
by (auto simp: ln-lower-1-def divide-simps)
```

4.1 Upper Bound 3

```
definition ln-upper-3 :: real ⇒ real
where ln-upper-3 ≡ λx. (x + 5)*(x - 1) / (2*(2*x + 1))

definition diff-delta-ln-upper-3 :: real ⇒ real
where diff-delta-ln-upper-3 ≡ λx. (x - 1)^3 / ((2*x + 1)^2 * x)

lemma d-delta-ln-upper-3: x > 0 ==>
((λx. ln-upper-3 x - ln x) has-field-derivative diff-delta-ln-upper-3 x) (at x)
unfolding ln-upper-3-def diff-delta-ln-upper-3-def
apply (intro derivative-eq-intros | simp)+
```

```
apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)
done
```

Strict inequalities also possible

lemma *ln-upper-3-pos*:

assumes $1 \leq x$ shows $\ln(x) \leq \ln\text{-upper-}3 x$

```
apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-3])
```

```
apply (auto simp: diff-delta-ln-upper-3-def ln-upper-3-def)
```

```
done
```

lemma *ln-upper-3-neg*:

assumes $0 < x$ and $x1: x \leq 1$ shows $\ln(x) \leq \ln\text{-upper-}3 x$

```
apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-3])
```

using assms

```
apply (auto simp: diff-delta-ln-upper-3-def divide-simps ln-upper-3-def)
```

```
done
```

theorem *ln-upper-3*: $0 < x \implies \ln(x) \leq \ln\text{-upper-}3 x$

by (metis le-less-linear less-eq-real-def ln-upper-3-neg ln-upper-3-pos)

definition *ln-lower-3* :: real \Rightarrow real

where $\ln\text{-lower-}3 \equiv \lambda x. - \ln\text{-upper-}3 (\text{inverse } x)$

corollary *ln-lower-3*: $0 < x \implies \ln\text{-lower-}3 x \leq \ln x$

unfolding *ln-lower-3-def*

by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-3)

theorem *ln-lower-3-eq*: $0 < x \implies \ln\text{-lower-}3 x = (1/2)*(1 + 5*x)*(x - 1) / (x*(2 + x))$

unfolding *ln-lower-3-def ln-upper-3-def*

by (simp add: divide-simps) algebra

4.2 Upper Bound 5

definition *ln-upper-5* :: real \Rightarrow real

where $\ln\text{-upper-}5 x \equiv (x^2 + 19*x + 10)*(x - 1) / (3*(3*x^2 + 6*x + 1))$

definition *diff-delta-ln-upper-5* :: real \Rightarrow real

where $\text{diff-delta-}ln\text{-upper-}5 \equiv \lambda x. (x - 1)^5 / ((3*x^2 + 6*x + 1)^2*x)$

lemma *d-delta-ln-upper-5*: $x > 0 \implies$

$((\lambda x. \ln\text{-upper-}5 x - \ln x) \text{ has-field-derivative } \text{diff-delta-}ln\text{-upper-}5 x) \text{ (at } x)$

unfolding *ln-upper-5-def diff-delta-*ln-upper-5-def**

apply (intro derivative-eq-intros | simp add: add-nonneg-eq-0-iff) +

apply (simp add: divide-simps add-nonneg-eq-0-iff, algebra)

done

lemma *ln-upper-5-pos*:

assumes $1 \leq x$ shows $\ln(x) \leq \ln\text{-upper-}5 x$

```

apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-5])
apply (auto simp: diff-delta-ln-upper-5-def ln-upper-5-def)
done

lemma ln-upper-5-neg:
assumes 0 < x and x1: x ≤ 1 shows ln(x) ≤ ln-upper-5 x
apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-5])
using assms
apply (auto simp: diff-delta-ln-upper-5-def divide-simps ln-upper-5-def mult-less-0-iff)
done

theorem ln-upper-5: 0 < x ==> ln(x) ≤ ln-upper-5 x
by (metis le-less-linear less-eq-real-def ln-upper-5-neg ln-upper-5-pos)

definition ln-lower-5 :: real => real
where ln-lower-5 ≡ λx. - ln-upper-5 (inverse x)

corollary ln-lower-5: 0 < x ==> ln-lower-5 x ≤ ln x
unfolding ln-lower-5-def
by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-5)

theorem ln-lower-5-eq: 0 < x ==
ln-lower-5 x = (1/3)*(10*x^2 + 19*x + 1)*(x - 1) / (x*(x^2 + 6*x + 3))
unfolding ln-lower-5-def ln-upper-5-def
by (simp add: zero-less-mult-iff add-pos-pos dual-order.strict-implies-not-eq divide-simps)
algebra

```

4.3 Upper Bound 7

```

definition ln-upper-7 :: real => real
where ln-upper-7 x ≡ (3*x^3 + 131*x^2 + 239*x + 47)*(x - 1) / (12*(4*x^3 + 18*x^2 + 12*x + 1))

```

```

definition diff-delta-ln-upper-7 :: real => real
where diff-delta-ln-upper-7 ≡ λx. (x - 1)^7 / ((4*x^3 + 18*x^2 + 12*x + 1)^2 * x)

```

```

lemma d-delta-ln-upper-7: x > 0 ==>
((λx. ln-upper-7 x - ln x) has-field-derivative diff-delta-ln-upper-7 x) (at x)
unfolding ln-upper-7-def diff-delta-ln-upper-7-def
apply (intro derivative-eq-intros | simp)+
apply auto
apply (auto simp: add-pos-pos dual-order.strict-implies-not-eq divide-simps, algebra)
done

```

```

lemma ln-upper-7-pos:
assumes 1 ≤ x shows ln(x) ≤ ln-upper-7 x

```

```

apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-7])
apply (auto simp: diff-delta-ln-upper-7-def ln-upper-7-def)
done

lemma ln-upper-7-neg:
  assumes "0 < x" and "x1: x ≤ 1" shows "ln(x) ≤ ln-upper-7 x"
apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-7])
using assms
apply (auto simp: diff-delta-ln-upper-7-def divide-simps ln-upper-7-def mult-less-0-iff)
done

theorem ln-upper-7: "0 < x" ⟹ "ln(x) ≤ ln-upper-7 x"
  by (metis le-less-linear less-eq-real-def ln-upper-7-neg ln-upper-7-pos)

definition ln-lower-7 :: real ⇒ real
  where "ln-lower-7 ≡ λx. - ln-upper-7 (inverse x)"

corollary ln-lower-7: "0 < x" ⟹ "ln-lower-7 x ≤ ln x"
  unfolding ln-lower-7-def
  by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-7)

theorem ln-lower-7-eq: "0 < x" ⟹
  "ln-lower-7 x = (1/12)*(47*x^3 + 239*x^2 + 131*x + 3)*(x - 1) / (x*(x^3 + 12*x^2 + 18*x + 4))"
  unfolding ln-lower-7-def ln-upper-7-def
  by (simp add: zero-less-mult-iff add-pos-pos dual-order.strict-implies-not-eq divide-simps)
algebra

```

4.4 Upper Bound 9

```

definition ln-upper-9 :: real ⇒ real
  where "ln-upper-9 x ≡ (6*x^4 + 481*x^3 + 1881*x^2 + 1281*x + 131)*(x - 1) / (30 * (5*x^4 + 40*x^3 + 60*x^2 + 20*x + 1))"

definition diff-delta-ln-upper-9 :: real ⇒ real
  where "diff-delta-ln-upper-9 ≡ λx. (x - 1)^9 / (((5*x^4 + 40*x^3 + 60*x^2 + 20*x + 1)^2) * x)"

```

```

lemma d-delta-ln-upper-9: "x > 0" ⟹
  "((λx. ln-upper-9 x - ln x) has-field-derivative diff-delta-ln-upper-9 x) (at x)"
unfolding ln-upper-9-def diff-delta-ln-upper-9-def
apply (intro derivative-eq-intros | simp) +
apply auto
apply (auto simp: add-pos-pos dual-order.strict-implies-not-eq divide-simps, algebra)
done

```

lemma *ln-upper-9-pos*:

assumes $1 \leq x$ shows $\ln(x) \leq \ln\text{-upper-9 } x$

apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-9])

apply (auto simp: diff-delta-ln-upper-9-def ln-upper-9-def)

done

lemma *ln-upper-9-neg*:

assumes $0 < x$ and $x1: x \leq 1$ shows $\ln(x) \leq \ln\text{-upper-9 } x$

apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-9])

using assms

apply (auto simp: diff-delta-ln-upper-9-def divide-simps ln-upper-9-def mult-less-0-iff)

done

theorem *ln-upper-9*: $0 < x \implies \ln(x) \leq \ln\text{-upper-9 } x$

by (metis le-less-linear less-eq-real-def ln-upper-9-neg ln-upper-9-pos)

definition *ln-lower-9* :: *real* \Rightarrow *real*

where $\ln\text{-lower-9} \equiv \lambda x. - \ln\text{-upper-9} (\text{inverse } x)$

corollary *ln-lower-9*: $0 < x \implies \ln\text{-lower-9 } x \leq \ln x$

unfoldng *ln-lower-9-def*

by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-9)

theorem *ln-lower-9-eq*: $0 < x \implies$

$\ln\text{-lower-9 } x = (1/30)*(6 + 481*x + 1881*x^2 + 1281*x^3 + 131*x^4)*(x - 1) / (x*(5 + 40*x + 60*x^2 + 20*x^3 + x^4))$

unfoldng *ln-lower-9-def ln-upper-9-def*

by (simp add: zero-less-mult-iff add-pos-pos dual-order.strict-implies-not-eq divide-simps)

algebra

4.5 Upper Bound 11

Extended bounds start here

definition *ln-upper-11* :: *real* \Rightarrow *real*

where $\ln\text{-upper-11 } x \equiv$

$(5*x^5 + 647*x^4 + 4397*x^3 + 6397*x^2 + 2272*x + 142) * (x - 1) / (30*(6*x^5 + 75*x^4 + 200*x^3 + 150*x^2 + 30*x + 1))$

definition *diff-delta-ln-upper-11* :: *real* \Rightarrow *real*

where $\text{diff-delta-ln-upper-11} \equiv \lambda x. (x - 1)^{11} / ((6*x^5 + 75*x^4 + 200*x^3 + 150*x^2 + 30*x + 1)^{12} * x)$

lemma *d-delta-ln-upper-11*: $x > 0 \implies$

$((\lambda x. \ln\text{-upper-11 } x - \ln x) \text{ has-field-derivative } \text{diff-delta-ln-upper-11 } x) \text{ (at } x\text{)}$

```

unfolding ln-upper-11-def diff-delta-ln-upper-11-def
apply (intro derivative-eq-intros | simp)+
apply auto
apply (auto simp: add-pos-pos dual-order.strict-implies-not-eq divide-simps, algebra)
done

lemma ln-upper-11-pos:
assumes  $1 \leq x$  shows  $\ln(x) \leq \ln\text{-upper-11 } x$ 
apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-11])
apply (auto simp: diff-delta-ln-upper-11-def ln-upper-11-def)
done

lemma ln-upper-11-neg:
assumes  $0 < x$  and  $x : x \leq 1$  shows  $\ln(x) \leq \ln\text{-upper-11 } x$ 
apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-11])
using assms
apply (auto simp: diff-delta-ln-upper-11-def divide-simps ln-upper-11-def mult-less-0-iff)
done

theorem ln-upper-11:  $0 < x \implies \ln(x) \leq \ln\text{-upper-11 } x$ 
by (metis le-less-linear less-eq-real-def ln-upper-11-neg ln-upper-11-pos)

definition ln-lower-11 :: real  $\Rightarrow$  real
where ln-lower-11  $\equiv \lambda x. -\ln\text{-upper-11 } (inverse x)$ 

corollary ln-lower-11:  $0 < x \implies \ln\text{-lower-11 } x \leq \ln x$ 
unfoldng ln-lower-11-def
by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-11)

theorem ln-lower-11-eq:  $0 < x \implies$ 

$$\ln\text{-lower-11 } x = (1/30)*(142*x^5 + 2272*x^4 + 6397*x^3 + 4397*x^2 + 647*x + 5)*(x - 1) / (x*(x^5 + 30*x^4 + 150*x^3 + 200*x^2 + 75*x + 6))$$

unfoldng ln-lower-11-def ln-upper-11-def
by (simp add: zero-less-mult-iff add-pos-pos dual-order.strict-implies-not-eq divide-simps)
algebra

```

4.6 Upper Bound 13

```

definition ln-upper-13 :: real  $\Rightarrow$  real
where ln-upper-13  $x \equiv (353 + 8389*x + 20149*x^4 + 50774*x^3 + 38524*x^2 + 1921*x^5 + 10*x^6) * (x - 1) / (70*(1 + 42*x + 525*x^4 + 700*x^3 + 315*x^2 + 126*x^5 + 7*x^6))$ 

definition diff-delta-ln-upper-13 :: real  $\Rightarrow$  real
where diff-delta-ln-upper-13  $\equiv \lambda x. (x - 1)^{13} /$ 

```

```

((1 + 42*x + 525*x^4 + 700*x^3 + 315*x^2 + 126*x^5 +
7*x^6)^2*x)

lemma d-delta-ln-upper-13:  $x > 0 \implies$ 
   $((\lambda x. \ln\text{-upper-}13 x - \ln x) \text{ has-field-derivative } \text{diff-delta}\text{-ln-upper-}13 x)$  (at  $x$ )
unfolding ln-upper-13-def diff-delta-ln-upper-13-def
apply (intro derivative-eq-intros | simp)+
apply auto
apply (auto simp: add-pos-pos dual-order.strict-implies-not-eq divide-simps, algebra)
done

lemma ln-upper-13-pos:
assumes  $1 \leq x$  shows  $\ln(x) \leq \ln\text{-upper-}13 x$ 
apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-13])
apply (auto simp: diff-delta-ln-upper-13-def ln-upper-13-def)
done

lemma ln-upper-13-neg:
assumes  $0 < x$  and  $x \leq 1$  shows  $\ln(x) \leq \ln\text{-upper-}13 x$ 
apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-13])
using assms
apply (auto simp: diff-delta-ln-upper-13-def divide-simps ln-upper-13-def mult-less-0-iff)
done

theorem ln-upper-13:  $0 < x \implies \ln(x) \leq \ln\text{-upper-}13 x$ 
by (metis le-less-linear less-eq-real-def ln-upper-13-neg ln-upper-13-pos)

definition ln-lower-13 :: real  $\Rightarrow$  real
where ln-lower-13  $\equiv \lambda x. -\ln\text{-upper-}13$  (inverse  $x$ )

corollary ln-lower-13:  $0 < x \implies \ln\text{-lower-}13 x \leq \ln x$ 
unfolding ln-lower-13-def
by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-13)

theorem ln-lower-13-eq:  $0 < x \implies$ 
 $\ln\text{-lower-}13 x = (1/70)*(10 + 1921*x + 20149*x^2 + 50774*x^3 + 38524*x^4$ 
 $+ 8389*x^5 + 353*x^6)*(x - 1) /$ 
 $(x*(7 + 126*x + 525*x^2 + 700*x^3 + 315*x^4 + 42*x^5 +$ 
 $x^6))$ 
unfolding ln-lower-13-def ln-upper-13-def
by (simp add: zero-less-mult-iff add-pos-pos dual-order.strict-implies-not-eq divide-simps)
algebra

```

4.7 Upper Bound 15

```

definition ln-upper-15 :: real  $\Rightarrow$  real
where ln-upper-15  $x \equiv$ 

```

```


$$(1487 + 49199*x + 547235*x^4 + 718735*x^3 + 334575*x^2 +$$


$$141123*x^5 + 35*x^7 + 9411*x^6)*(x - 1) /$$


$$(280*(1 + 56*x + 2450*x^4 + 1960*x^3 + 588*x^2 + 1176*x^5 +$$


$$8*x^7 + 196*x^6))$$


definition diff-delta-ln-upper-15 :: real  $\Rightarrow$  real
where diff-delta-ln-upper-15
 $\equiv \lambda x. (x - 1)^{15} / ((1 + 56*x + 2450*x^4 + 1960*x^3 + 588*x^2 + 8*x^7 + 196*x^6 + 1176*x^5)^2 * x)$ 

lemma d-delta-ln-upper-15:  $x > 0 \implies$ 
 $((\lambda x. \ln\text{-upper-15 } x - \ln x) \text{ has-field-derivative } \text{diff-delta-ln-upper-15 } x) \text{ (at } x)$ 
unfolding ln-upper-15-def diff-delta-ln-upper-15-def
apply (intro derivative-eq-intros | simp)+
apply auto
apply (auto simp: add-pos-pos dual-order.strict-implies-not-eq divide-simps, algebra)
done

lemma ln-upper-15-pos:
assumes  $1 \leq x$  shows  $\ln(x) \leq \ln\text{-upper-15 } x$ 
apply (rule gen-upper-bound-increasing [OF assms d-delta-ln-upper-15])
apply (auto simp: diff-delta-ln-upper-15-def ln-upper-15-def)
done

lemma ln-upper-15-neg:
assumes  $0 < x$  and  $x1: x \leq 1$  shows  $\ln(x) \leq \ln\text{-upper-15 } x$ 
apply (rule gen-upper-bound-decreasing [OF x1 d-delta-ln-upper-15])
using assms
apply (auto simp: diff-delta-ln-upper-15-def divide-simps ln-upper-15-def mult-less-0-iff)
done

theorem ln-upper-15:  $0 < x \implies \ln(x) \leq \ln\text{-upper-15 } x$ 
by (metis le-less-linear less-eq-real-def ln-upper-15-neg ln-upper-15-pos)

definition ln-lower-15 :: real  $\Rightarrow$  real
where ln-lower-15  $\equiv \lambda x. -\ln\text{-upper-15 } (inverse x)$ 

corollary ln-lower-15:  $0 < x \implies \ln\text{-lower-15 } x \leq \ln x$ 
unfolding ln-lower-15-def
by (metis ln-inverse inverse-positive-iff-positive minus-le-iff ln-upper-15)

theorem ln-lower-15-eq:  $0 < x \implies$ 
 $\ln\text{-lower-15 } x = (1/280)*(35 + 9411*x + 141123*x^2 + 547235*x^3 +$ 
 $718735*x^4 + 334575*x^5 + 49199*x^6 + 1487*x^7)*(x - 1) /$ 
 $(x*(8 + 196*x + 1176*x^2 + 2450*x^3 + 1960*x^4 + 588*x^5 +$ 
 $56*x^6 + x^7))$ 
unfolding ln-lower-15-def ln-upper-15-def

```

by (*simp add: zero-less-mult-iff add-pos-pos dual-order.strict-implies-not-eq divide-simps*) *algebra*

end

Chapter 5

Sine and Cosine Upper and Lower Bounds

```
theory Sin-Cos-Bounds
imports Bounds-Lemmas
```

```
begin
```

5.1 Simple base cases

Upper bound for $(0 :: 'a) \leq x$

```
lemma sin-le-arg:
```

```
  fixes x :: real
```

```
  shows  $0 \leq x \implies \sin x \leq x$ 
```

```
  by (fact sin-x-le-x)
```

```
lemma cos-ge-1-arg:
```

```
  fixes x :: real
```

```
  assumes  $0 \leq x$ 
```

```
  shows  $1 - x \leq \cos x$ 
```

```
  apply (rule gen-lower-bound-increasing [OF assms])
```

```
  apply (intro derivative-eq-intros, auto)
```

```
  done
```

```
lemmas sin-Taylor-0-upper-bound-pos = sin-le-arg — MetiTarski bound
```

```
lemma cos-Taylor-1-lower-bound:
```

```
  fixes x :: real
```

```
  assumes  $0 \leq x$ 
```

```
  shows  $(1 - x^2 / 2) \leq \cos x$ 
```

```
  apply (rule gen-lower-bound-increasing [OF assms])
```

```
  apply (intro derivative-eq-intros)
```

```
  apply (rule refl | simp add: sin-le-arg)+
```

```
  done
```

```

lemma sin-Taylor-1-lower-bound:
  fixes x :: real
  assumes 0 ≤ x
  shows (x - x ^ 3 / 6) ≤ sin x
  apply (rule gen-lower-bound-increasing [OF assms])
  apply (intro derivative-eq-intros)
  apply (rule refl | simp add: cos-Taylor-1-lower-bound) +
  done

```

5.2 Taylor series approximants

```

definition sinpoly :: [nat,real] ⇒ real
  where sinpoly n = (λx. ∑ k< n. sin-coeff k * x ^ k)

```

```

definition cospoly :: [nat,real] ⇒ real
  where cospoly n = (λx. ∑ k< n. cos-coeff k * x ^ k)

```

```

lemma sinpoly-Suc: sinpoly (Suc n) = (λx. sinpoly n x + sin-coeff n * x ^ n)
  by (simp add: sinpoly-def)

```

```

lemma cospoly-Suc: cospoly (Suc n) = (λx. cospoly n x + cos-coeff n * x ^ n)
  by (simp add: cospoly-def)

```

```

lemma sinpoly-minus [simp]: sinpoly n (-x) = - sinpoly n x
  by (induct n) (auto simp: sinpoly-def sin-coeff-def)

```

```

lemma cospoly-minus [simp]: cospoly n (-x) = cospoly n x
  by (induct n) (auto simp: cospoly-def cos-coeff-def)

```

```

lemma d-sinpoly-cospoly:
  (sinpoly (Suc n) has-field-derivative cospoly n x) (at x)
  proof (induction n)
    case 0 show ?case
      by (simp add: sinpoly-def cospoly-def)
    next
    case (Suc n) show ?case
      proof (cases n=0)
        case True then show ?thesis
        by (simp add: sinpoly-def sin-coeff-def cospoly-def)
      next
        case False then
        have xn: x ^ (n - Suc 0) * x = x ^ n
        by (metis Suc-pred mult.commute not-gr0 power-Suc)
        show ?thesis using Suc False
        apply (simp add: sinpoly-Suc [of Suc n] cospoly-def)
        apply (intro derivative-eq-intros | simp) +

```

```

apply (simp add: xn mult.assoc sin-coeff-def cos-coeff-def divide-simps del:
fact-Suc)
apply (simp add: algebra-simps)
done
qed
qed

lemma d-cospoly-sinpoly:
  (cospoly (Suc n) has-field-derivative -sinpoly n x) (at x)
proof (induction n)
  case 0 show ?case
    by (simp add: sinpoly-def cospoly-def)
next
  case (Suc n) show ?case
  proof (cases n=0)
    case True then show ?thesis
    by (simp add: sinpoly-def cospoly-def cos-coeff-def)
  next
    case False then
      have xn:  $x^{\wedge}(n - Suc 0) * x = x^{\wedge}n$ 
      by (metis Suc-pred mult.commute not-gr0 power-Suc)
      have m1: odd n  $\implies (-1 :: real)^{\wedge}((n - Suc 0) \text{ div } 2) = -((-1)^{\wedge}(Suc n \text{ div } 2))$ 
      by (cases n) simp-all
      show ?thesis using Suc False
        apply (simp add: cospoly-Suc [of Suc n] sinpoly-def)
        apply (intro derivative-eq-intros | simp)+
        apply (simp add: xn mult.assoc sin-coeff-def cos-coeff-def m1 divide-simps del:
fact-Suc)
        apply (simp add: algebra-simps)
        done
      qed
    qed
  qed

```

5.3 Inductive proof of sine inequalities

```

lemma sinpoly-lb-imp-cospoly-ub:
  assumes x0:  $0 \leq x$  and k0:  $k > 0$  and  $\bigwedge x. 0 \leq x \implies \text{sinpoly} (k - 1) x \leq \sin x$ 
  shows  $\cos x \leq \text{cospoly} k x$ 
  apply (rule gen-lower-bound-increasing [OF x0])
  apply (intro derivative-eq-intros | simp)+
  using d-cospoly-sinpoly [of k - 1] assms
  apply auto
  apply (simp add: cospoly-def)
  done

lemma cospoly-ub-imp-sinpoly-ub:
  assumes x0:  $0 \leq x$  and k0:  $k > 0$  and  $\bigwedge x. 0 \leq x \implies \cos x \leq \text{cospoly} (k - 1) x$ 
  shows  $\sin x \leq \text{sinpoly} k x$ 

```

```

apply (rule gen-lower-bound-increasing [OF x0])
apply (intro derivative-eq-intros | simp)+
using d-sinpoly-cospoly [of k - 1] assms
apply auto
apply (simp add: sinpoly-def)
done

lemma sinpoly-ub-imp-cospoly-lb:
assumes x0:  $0 \leq x$  and k0:  $k > 0$  and  $\bigwedge x. 0 \leq x \implies \sin x \leq \text{sinpoly}(k - 1) x$ 
shows cospoly k x  $\leq \cos x$ 
apply (rule gen-lower-bound-increasing [OF x0])
apply (intro derivative-eq-intros | simp)+
using d-cospoly-sinpoly [of k - 1] assms
apply auto
apply (simp add: cospoly-def)
done

lemma cospoly-lb-imp-sinpoly-lb:
assumes x0:  $0 \leq x$  and k0:  $k > 0$  and  $\bigwedge x. 0 \leq x \implies \cos x \leq \text{cospoly}(k - 1) x \leq \sin x$ 
shows sinpoly k x  $\leq \sin x$ 
apply (rule gen-lower-bound-increasing [OF x0])
apply (intro derivative-eq-intros | simp)+
using d-sinpoly-cospoly [of k - 1] assms
apply auto
apply (simp add: sinpoly-def)
done

lemma
assumes  $0 \leq x$ 
shows sinpoly-lower-nonneg:  $\text{sinpoly}(4 * \text{Suc } n) x \leq \sin x$  (is ?th1)
and sinpoly-upper-nonneg:  $\sin x \leq \text{sinpoly}(\text{Suc}(\text{Suc}(4*n))) x$  (is ?th2)
proof -
have sinpoly (4 * Suc n) x  $\leq \sin x \wedge \sin x \leq \text{sinpoly}(\text{Suc}(\text{Suc}(4*n))) x$ 
using assms
apply (induction n arbitrary: x)
apply (simp add: sinpoly-def sin-coeff-def sin-Taylor-1-lower-bound sin-Taylor-0-upper-bound-pos
lessThan-nat-numeral fact-numeral)
apply (auto simp: cospoly-lb-imp-sinpoly-lb sinpoly-ub-imp-cospoly-lb cospoly-ub-imp-sinpoly-ub
sinpoly-lb-imp-cospoly-ub)
done
then show ?th1 ?th2
using assms
by auto
qed

```

5.4 Collecting the results

corollary sinpoly-upper-nonpos:
 $x \leq 0 \implies \sin x \leq \text{sinpoly}(4 * \text{Suc } n) x$

```

using sinpoly-lower-nonneg [of  $-x$   $n$ ]
by simp

corollary sinpoly-lower-nonpos:
   $x \leq 0 \implies \text{sinpoly}(\text{Suc}(\text{Suc}(\text{Suc}(4*n)))) x \leq \sin x$ 
using sinpoly-upper-nonneg [of  $-x$   $n$ ]
by simp

corollary cospoly-lower-nonneg:
   $0 \leq x \implies \text{cospoly}(\text{Suc}(\text{Suc}(\text{Suc}(4*n)))) x \leq \cos x$ 
by (auto simp: sinpoly-upper-nonneg sinpoly-ub-imp-copoly-lb)

lemma cospoly-lower:
   $\text{cospoly}(\text{Suc}(\text{Suc}(\text{Suc}(4*n)))) x \leq \cos x$ 
proof (cases rule: le-cases [of  $0$   $x$ ])
  case le then show ?thesis
    by (simp add: cospoly-lower-nonneg)
next
  case ge then show ?thesis using cospoly-lower-nonneg [of  $-x$ ]
    by simp
qed

lemma cospoly-upper-nonneg:
  assumes  $0 \leq x$ 
  shows  $\cos x \leq \text{cospoly}(\text{Suc}(4*n)) x$ 
proof (cases  $n$ )
  case 0 then show ?thesis
    by (simp add: cospoly-def)
next
  case ( $\text{Suc } m$ )
  then show ?thesis
    using sinpoly-lower-nonneg [of  $-m$ ] assms
    by (auto simp: sinpoly-lb-imp-copoly-ub)
qed

lemma cospoly-upper:
   $\cos x \leq \text{cospoly}(\text{Suc}(4*n)) x$ 
proof (cases rule: le-cases [of  $0$   $x$ ])
  case le then show ?thesis
    by (simp add: cospoly-upper-nonneg)
next
  case ge then show ?thesis using cospoly-upper-nonneg [of  $-x$ ]
    by simp
qed

end

```

Chapter 6

Square Root Upper and Lower Bounds

```
theory Sqrt-Bounds
imports Bounds-Lemmas
HOL-Library.Sum-of-Squares

begin

Covers all bounds used in sqrt-upper.ax, sqrt-lower.ax and sqrt-extended.ax.

6.1 Upper bounds

primrec sqrtu :: [real,nat] ⇒ real where
  sqrtu x 0 = (x+1)/2
| sqrtu x (Suc n) = (sqrtu x n + x/sqrtu x n) / 2

lemma sqrtu-upper: x ≤ sqrtu x n ^ 2
proof (induction n)
  case 0 show ?case
    apply (simp add: power2-eq-square)
    apply (sos (((A<0 * R<1) + (R<1 * (R<1 * [~1*x + 1]^2))))) done
  next
  case (Suc n)
    have xy: ∀y. [x ≤ y * y; y ≠ 0] ⇒ x * (2 * (y * y)) ≤ x * x + y * (y * y)
      by (sos ((((A<0 * A<1) * R<1) + ((A<0 * R<1) * (R<1 * [~1*y^2 + x]^2)))) &
        (((((A<0 * A<1) * R<1) + ((A<0 * R<1) * (R<1 * [~1*y^2 + x]^2))))) show ?case using Suc
      by (auto simp: power2-eq-square algebra-simps divide-simps xy)
qed
```

```

lemma sqrtu-numeral:
  sqrtu x (numeral n) = (sqrtu x (pred-numeral n) + x/sqrtu x (pred-numeral n))
  / 2
  by (simp add: numeral-eq-Suc)

lemma sqrtu-gt-0:  $x \geq 0 \implies \sqrt{u} x n > 0$ 
  apply (induct n)
  apply (auto simp: field-simps)
  by (metis add-strict-increasing2 mult-zero-left not-real-square-gt-zero)

theorem gen-sqrt-upper:  $0 \leq x \implies \sqrt{x} \leq \sqrt{u} x n$ 
  using real-sqrt-le-mono [OF sqrtu-upper [of x n]]
  by auto (metis abs-of-nonneg dual-order.strict-iff-order sqrtu-gt-0)

lemma sqrt-upper-bound-0:
  assumes  $x \geq 0$  shows  $\sqrt{x} \leq (x+1)/2$  (is -  $\leq ?rhs$ )
  proof -
    have  $\sqrt{x} \leq \sqrt{u} x 0$ 
    by (metis assms gen-sqrt-upper)
    also have ... = ?rhs
    by (simp add: divide-simps)
    finally show ?thesis .
  qed

lemma sqrt-upper-bound-1:
  assumes  $x \geq 0$  shows  $\sqrt{x} \leq (1/4)*(x^2+6*x+1) / (x+1)$  (is -  $\leq ?rhs$ )
  proof -
    have  $\sqrt{x} \leq \sqrt{u} x 1$ 
    by (metis assms gen-sqrt-upper)
    also have ... = ?rhs
    by (simp add: divide-simps) algebra
    finally show ?thesis .
  qed

lemma sqrtu-2-eq:
  
$$\sqrt{u} x 2 = (1/8)*(x^4 + 28*x^3 + 70*x^2 + 28*x + 1) / ((x + 1)*(x^2 + 6*x + 1))$$

  by (simp add: sqrtu-numeral divide-simps) algebra

lemma sqrt-upper-bound-2:
  assumes  $x \geq 0$ 
  shows  $\sqrt{x} \leq (1/8)*(x^4 + 28*x^3 + 70*x^2 + 28*x + 1) / ((x + 1)*(x^2 + 6*x + 1))$ 
  by (metis assms gen-sqrt-upper sqrtu-2-eq)

lemma sqrtu-4-eq:
  
$$x \geq 0 \implies \sqrt{u} x 4 = (1/32)*(225792840*x^6 + 64512240*x^5 + 601080390*x^8 + 471435600*x^7 + 496*x + 1 + 35960*x^2 + 120*x^5 + 8008*x^3 + 70*x^2 + 28*x + 1) / ((x + 1)*(x^2 + 6*x + 1)*(x^4 + 28*x^3 + 70*x^2 + 28*x + 1)*(1820*x^6 + 8008*x^5 + x^8 + 120*x^7 + 35960*x^4 + 471435600*x^3 + 601080390*x^2 + 64512240*x + 225792840))$$


```

by (*simp add: sqrtu-numeral divide-simps add-nonneg-eq-0-iff*) *algebra*

lemma *sqrt-upper-bound-4*:

assumes $x \geq 0$

shows $\sqrt{x} \leq (1/32)*(225792840*x^6 + 64512240*x^5 + 601080390*x^8 + 471435600*x^7 + 496*x + 1 + 35)$
 $/ ((x+1)*(x^2+6*x+1)*(x^4+28*x^3+70*x^2+28*x+1)*(1820*x^6+8008*x^5+x^8+120*x^7+1))$

by (*metis assms gen-sqrt-upper sqrtu-4-eq*)

lemma *gen-sqrt-upper-scaled*:

assumes $0 \leq x \ 0 < u$

shows $\sqrt{x} \leq \sqrt{u} (x*u^2) n / u$

proof -

have $\sqrt{x} = \sqrt{x} * \sqrt{u^2} / u$

using *assms*

by *simp*

also have ... $= \sqrt{(x*u^2)} / u$

by (*metis real-sqrt-mult*)

also have ... $\leq \sqrt{u} (x*u^2) n / u$

using *assms*

by (*simp add: divide-simps*) (*metis gen-sqrt-upper zero-le-mult-iff zero-le-power2*)

finally show ?thesis .

qed

lemma *sqrt-upper-bound-2-small*:

assumes $0 \leq x$

shows $\sqrt{x} \leq (1/32)*(65536*x^4 + 114688*x^3 + 17920*x^2 + 448*x + 1) / ((16*x + 1)*(256*x^2 + 96*x + 1))$

apply (*rule order-trans [OF gen-sqrt-upper-scaled [of x 4 2] eq-refl]*)

using *assms*

apply (*auto simp: sqrtu-2-eq*)

apply (*simp add: divide-simps*)

apply *algebra*

done

lemma *sqrt-upper-bound-2-large*:

assumes $0 \leq x$

shows $\sqrt{x} \leq (1/32)*(65536 + 114688*x + 17920*x^2 + 448*x^3 + x^4) / ((x + 16)*(256 + 96*x + x^2))$

apply (*rule order-trans [OF gen-sqrt-upper-scaled [of x 1/4 2] eq-refl]*)

using *assms*

apply (*auto simp: sqrtu-2-eq*)

apply (*simp add: divide-simps*)

apply *algebra*

done

6.2 Lower bounds

lemma *sqrt-lower-bound-id*:

assumes $0 \leq x \ x \leq 1$

```

shows  $x \leq \sqrt{x}$ 
proof –
  have  $x^2 \leq x$  using assms
    by (metis one-le-numeral power-decreasing power-one-right)
  then show ?thesis
    by (metis real-le-rsqrt)
qed

definition sqrtl :: [real,nat] ⇒ real where
  sqrtl x n = x/sqrtn x n

lemma sqrtl-lower:  $0 \leq x \implies \text{sqrtl } x \leq \sqrt{x}$ 
  unfolding sqrtl-def using sqrtu-upper [of x n]
  by (auto simp: power2-eq-square divide-simps mult-left-mono)

theorem gen-sqrt-lower:  $0 \leq x \implies \text{sqrtl } x \leq \sqrt{x}$ 
  using real-sqrt-le-mono [OF sqrtl-lower [of x n]]
  by auto

lemma sqrt-lower-bound-0:
  assumes  $x \geq 0$  shows  $2*x/(x+1) \leq \sqrt{x}$  (is ?lhs ≤ -)
proof –
  have ?lhs = sqrtl x 0
    by (simp add: sqrtl-def)
  also have ... ≤ sqrt x
    by (metis assms gen-sqrt-lower)
  finally show ?thesis .
qed

lemma sqrt-lower-bound-1:
  assumes  $x \geq 0$  shows  $4*x*(x+1) / (x^2 + 6*x + 1) \leq \sqrt{x}$  (is ?lhs ≤ -)
proof –
  have ?lhs = sqrtl x 1 using assms
    by (simp add: sqrtl-def power-eq-if algebra-simps divide-simps)
  also have ... ≤ sqrt x
    by (metis assms gen-sqrt-lower)
  finally show ?thesis .
qed

lemma sqrtl-2-eq:  $\text{sqrtl } x^2 = 8*x*(x+1)*(x^2 + 6*x + 1) / (x^4 + 28*x^3 + 70*x^2 + 28*x + 1)$ 
  using sqrtu-gt-0 [of x 2]
  by (simp add: sqrtl-def sqrtu-2-eq)

lemma sqrt-lower-bound-2:
  assumes  $x \geq 0$ 
  shows  $8*x*(x+1)*(x^2 + 6*x + 1) / (x^4 + 28*x^3 + 70*x^2 + 28*x + 1) \leq \sqrt{x}$ 
  by (metis assms sqrtl-2-eq gen-sqrt-lower)

```

```

lemma sqrtl-4-eq:  $x \geq 0 \implies$ 
  sqrtl  $x^4$ 
   $= (32*x*(x+1)*(x^2+6*x+1)*(x^4+28*x^3+70*x^2+28*x+1)*(1820*x^6+8008*x^5+x^8+120*x^7+$ 
   $/ (225792840*x^6+64512240*x^5+601080390*x^8+471435600*x^7+496*x+1+35960*x^2+906192*x^3+$ 
  using sqrtu-gt-0 [of  $x^4$ ]
  by (simp add: sqrtl-def sqrtu-4-eq)

lemma sqrt-lower-bound-4:
  assumes  $x \geq 0$ 
  shows  $(32*x*(x+1)*(x^2+6*x+1)*(x^4+28*x^3+70*x^2+28*x+1)*(1820*x^6+8008*x^5+x^8+120*x^7+$ 
   $/ (225792840*x^6+64512240*x^5+601080390*x^8+471435600*x^7+496*x+1+35960*x^2+906192*x^3+$ 
   $\leq \sqrt{x}$ 
  by (metis assmss sqrtl-4-eq gen-sqrt-lower)

lemma gen-sqrt-lower-scaled:
  assumes  $0 \leq x \quad 0 < u$ 
  shows sqrtl  $(x*u^2) / u \leq \sqrt{x} / u$ 
  proof -
    have sqrtl  $(x*u^2) / u \leq \sqrt{(x*u^2) / u}$ 
    using assmss
    by (simp add: divide-simps) (metis gen-sqrt-lower zero-le-mult-iff zero-le-power2)
    also have ... =  $\sqrt{x} * \sqrt{u^2} / u$ 
    by (metis real-sqrt-mult)
    also have ... =  $\sqrt{x}$ 
    using assmss
    by simp
    finally show ?thesis .
  qed

lemma sqrt-lower-bound-2-small:
  assumes  $0 \leq x$ 
  shows  $32*x*(16*x + 1)*(256*x^2 + 96*x + 1) / (65536*x^4 + 114688*x^3$ 
   $+ 17920*x^2 + 448*x + 1) \leq \sqrt{x}$ 
  apply (rule order-trans [OF eq-refl gen-sqrt-lower-scaled [of  $x^4 2$ ]])
  using assmss
  apply (auto simp: sqrtl-2-eq)
  apply (simp add: divide-simps)
  apply algebra
  done

lemma sqrt-lower-bound-2-large:
  assumes  $0 \leq x$ 
  shows  $32*x*(x + 16)*(x^2 + 96*x + 256) / (x^4 + 448*x^3 + 17920*x^2$ 
   $+ 114688*x + 65536) \leq \sqrt{x}$ 
  apply (rule order-trans [OF eq-refl gen-sqrt-lower-scaled [of  $x^4 2$ ]])
  using assmss
  apply (auto simp: sqrtl-2-eq)
  apply (simp add: divide-simps)

```

done

end

Bibliography

- [1] B. Akbarpour and L. Paulson. MetiTarski: An automatic theorem prover for real-valued special functions. 44(3):175–205, Mar. 2010.