Wooley's Discrete Inequality

Angeliki Koutsoukou-Argyraki
Royal Holloway, University of London and University of Cambridge

February 6, 2026

Abstract

This is a formalisation of the proof of an inequality by Trevor D. Wooley attesting that when $\lambda > 0$,

$$\min_{r \in \mathbb{N}}(r + \lambda/r) \leq \sqrt{4\lambda + 1}$$

with equality if and only if $\lambda = m(m-1)$ for some positive integer m.

Contents

1 Wooley's Discrete Inequality **1**
1.1 General elementary technical lemmas 2
1.2 Trivial case, where we minimise over all positive real values of r 2
1.3 Main result: Inequality for the discrete version 2
1.4 Special case: Equality for the discrete version 2

1 Wooley's Discrete Inequality

theory *Wooley-Elementary-Discrete-Inequality*
imports *HOL−Library.Quadratic-Discriminant HOL−Real-Asymp.Real-Asymp*

begin

This is a formalisation of the proof of an inequality by Trevor D. Wooley attesting that when $\lambda > 0$,

$$\min_{r \in \mathbb{N}}(r + \lambda/r) \leq \sqrt{4\lambda + 1}$$

with equality if and only if $\lambda = m(m-1)$ for some positive integer m. The source is the note "An Elementary Discrete Inequality" available on Wooley's webpage [1]: https://www.math.purdue.edu/~twooley/publ/20230410discineq.pdf.

1.1 General elementary technical lemmas

lemma *obtains-nat-in-interval*:
fixes x::*real* **assumes** $x \geq 0$
obtains c::*nat* **where** $c \in \{x<..\ x+1\}$
⟨*proof*⟩

lemma *obtains-nat-in-interval-greater-leq*:
fixes x::*real* **assumes** $x \geq 0$
obtains c::*nat* **where** $c > x$ **and** $c \leq x+1$
⟨*proof*⟩

lemma *obtains-nat-in-interval-half*:
fixes x::*real* **assumes** $x \geq 1/2$
obtains c::*nat* **where** $c > x - (1/2)$ **and** $c \leq x+1/2$
⟨*proof*⟩

1.2 Trivial case, where we minimise over all positive real values of r

theorem *elementary-ineq-Wooley-real*:
fixes l::*real* **and** g::*real* ⇒ *real*
assumes $l>0$ **and** $\forall\ r \in R.\ g\ r = r+(l/r)$
and $R=\{r$::*real*. $r>0\}$
shows $(\forall\ r \in R.\ g\ r \geq\ 2*\ sqrt(l)) \wedge (\forall\ r \in R.\ g\ (sqrt(l)) \leq g\ r)$
⟨*proof*⟩

1.3 Main result: Inequality for the discrete version

theorem *elementary-discrete-ineq-Wooley*:
fixes l::*real* **and** g::*nat* ⇒ *real*
assumes $l>0$ **and** $R = \{r$::*nat*. $r>0\}$ **and** $\forall\ r \in R.\ g\ r = r+\ (l/r)$
shows $(INF\ r \in R.\ g\ r) \leq sqrt(4*l+1)$

⟨*proof*⟩

1.4 Special case: Equality for the discrete version

We will now show a special case of the main result where equality holds instead of inequality.

We will need to make use of the following technical lemma, which will be used so as to guarantee that there exists a $p \in R$ for which the INF of $g(r)$ equals to $g(p)$. To this end, we will show that here the infimum INF can be identified with the minimum Min by restricting to a finite set. As the operator Min in Isabelle is used for finite sets and R is infinite, we used INF in the original formulation, however here Min and INF can be identified.

The following technical lemma is by Larry Paulson:

lemma *restrict-to-min*:

fixes l::*real* **and** g::$nat \Rightarrow$ *real*
assumes $l>0$ **and** *R-def*: $R=\{r::nat.\ r>0\}$ **and** *g-def*: $\forall\ r.\ g\ r = r + (l/r)$
obtains F **where** *finite* F $F \subseteq R$ $(INF\ r \in R.\ g\ r) = Min\ (g\ `\ F)$ $F \neq \{\}$

⟨*proof*⟩

We will make use of the following calculation, which is convenient to formulate separately as a lemma.

lemma *elementary-discrete-ineq-Wooley-quadratic-eq-sol*:
fixes l::*real* **and** g::$nat \Rightarrow$ *real*
assumes $l>0$ **and** $\forall\ r.\ g\ r = r + (l/r)$ **and** $g\ r = sqrt(4*l+1)$
shows $(r = 1/2 + (1/2)*\ sqrt(\ 4*l\ +1)) \vee (r = -\ 1/2 + (1/2)*\ sqrt(4*l\ +1))$
⟨*proof*⟩

The special case with equality involves a double implication (iff), and we start by showing one direction.

theorem *elementary-discrete-ineq-Wooley-special-case-1*:
fixes l::*real* **and** g::$nat \Rightarrow$ *real* **assumes** $l>0$ **and** $R=\{r::nat.\ r>0\}$ **and** $\forall\ r.\ g\ r = r + (l/r)$
and $(INF\ r \in R.\ g\ r) = sqrt(4*l+1)$
shows $\exists\ m::nat.\ l = m*(m-1)$

⟨*proof*⟩

Now we show the other direction.

theorem *elementary-discrete-ineq-Wooley-special-case-2*:
fixes l::*real* **and** g::$nat \Rightarrow$ *real*
assumes $l>0$ **and** $R=\{r::nat.\ r>0\}$ **and** $\forall\ r.\ g\ r = r + (l/r)$ **and** $\exists\ m::nat.\ l = m*(m-1)$
shows $(INF\ r \in R.\ g\ r) = sqrt(4*l+1)$

⟨*proof*⟩

Finally, for convenience and completeness, we state the special case where equality holds formulated with the double implication and moreover including the values for which the INF (i.e. minimum here as we have seen) is attained as previously calculated.

theorem *elementary-discrete-ineq-Wooley-special-case-iff*:
fixes l::*real* **and** g::$nat \Rightarrow$ *real*
assumes $l>0$ **and** $R=\{r::nat.\ r>0\}$ **and** $\forall\ r.\ g\ r = r + (l/r)$
shows $((INF\ r \in R.\ g\ r) = sqrt(4*l+1)) \longleftrightarrow (\exists\ m::nat.\ l = m*(m-1))$
and
$g\ p = sqrt(4*l+1) \longrightarrow (p = 1/2 + (1/2)*\ sqrt(\ 4*l\ +1)) \vee (p = -1/2 + (1/2)*\ sqrt(4*l\ +1))$
⟨*proof*⟩

end

References

[1] T. D. Wooley. An elementary discrete inequality. https://www.math.purdue.edu/~twooley/publ/20230410discineq.pdf.